

$$1. \int_0^{\pi/4} \frac{\sin x + \cos x}{9+16\sin 2x} dx$$

sol: let

$$\sin x - \cos x = t$$

diff . w.r.t 'x'

$$(\cos x + \sin x)dx = dt$$

$$L.L: x = 0 \Rightarrow t = \sin 0 - \cos 0 = 0 - 1 = -1$$

$$U.L: x = \frac{\pi}{4} \Rightarrow t = \sin\left(\frac{\pi}{4}\right) - \cos\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = 0$$

$$\sin x - \cos x = t \quad S.O.B$$

$$\Rightarrow (\sin x - \cos x)^2 = t^2$$

$$\Rightarrow \sin^2 x + \cos^2 x - 2\sin x \cos x = t^2$$

$$\Rightarrow 1 - \sin 2x = t^2$$

$$1 - t^2 = \sin 2x$$

$$I = \int_0^{\pi/4} \frac{\sin x + \cos x}{9+16\sin 2x} dx$$

$$= \int_{-1}^0 \frac{1}{9+16(1-t^2)} dt$$

$$= \int_{-1}^0 \frac{1}{9+16-16t^2} dt$$

$$= \int_{-1}^0 \frac{1}{25-16t^2} dt$$

$$= \int_{-1}^0 \frac{1}{(5-4t)(5+4t)} dt$$

$$\therefore \int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + c$$

$$= \frac{\left[\frac{1}{2[5]} \log \left| \frac{5+4t}{4-5t} \right| \right]_0^{-1}}{4}$$

$$= \frac{1}{40} \left[\log \left| \frac{5+0}{5-0} \right| - \log \left| \frac{5-4}{5+4} \right| \right]$$

$$= \frac{1}{40} \left[\log 1 - \log \frac{1}{9} \right]$$

$$\log 1 = 0$$

$$= \frac{1}{40} [0 - \log 3^{-2}] = \frac{1}{40} [2 \log 3]$$

$$= \frac{1}{20} \log 3$$

Laq Q.No 23

$$2. \int_0^1 \frac{\log(1+x)}{(1+x^2)} dx$$

sol: let

$$x = \tan \theta$$

diff . w.r.t 'x'

$$dx = \sec^2 \theta d\theta$$

$$L.L: x = 0 \Rightarrow \theta = 0$$

$$U.L: x = 1 \Rightarrow \theta = \frac{\pi}{4}$$

$$\int_0^1 \frac{\log(1+x)}{(1+x^2)} dx$$

$$= \int_0^{\pi/4} \frac{\log(1+\tan\theta)}{(1+\tan^2\theta)} \sec^2 \theta d\theta$$

$$I = \int_0^{\pi/4} \log(1 + \tan\theta) d\theta \dots \dots \dots (1)$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^{\pi/4} \log \left(1 + \tan \left[\frac{\pi}{4} - \theta \right] \right) d\theta$$

$$I = \int_0^{\pi/4} \log \left(1 + \frac{1-\tan\theta}{1+\tan\theta} \right) d\theta$$

$$I = \int_0^{\pi/4} \log \left(\frac{1+\tan\theta+1-\tan\theta}{1+\tan\theta} \right) d\theta$$

$$I = \int_0^{\pi/4} \log \left(\frac{2}{1+\tan\theta} \right) d\theta$$

$$\log(a/b) = \log a - \log b$$

$$I = \int_0^{\pi/4} \log 2 d\theta - \int_0^{\pi/4} \log(1 + \tan\theta) d\theta$$

$$I + I = \int_0^{\pi/4} \log 2 d\theta$$

$$2I = \log 2 \int_0^{\pi/4} (1) d\theta$$

$$2I = \log 2 [\theta]_0^{\pi/4}$$

$$2I = \log 2 \left[\frac{\pi}{4} - 0 \right]$$

$$I = \frac{\pi}{8} \log 2$$

Definite integrals

3. $\int_0^\pi \frac{x \cdot \sin x}{1 + \sin x} dx$

Sol:

$$I = \int_0^\pi \frac{x \cdot \sin x}{1 + \sin x} dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^\pi \frac{(\pi-x) \cdot \sin(\pi-x)}{1 + \sin(\pi-x)} dx$$

$$I = \int_0^\pi \frac{(\pi-x) \cdot \sin x}{1 + \sin x} dx$$

$$I = \int_0^\pi \frac{\pi \sin x}{1 + \sin x} dx - \int_0^\pi \frac{x \cdot \sin x}{1 + \sin x} dx$$

$$I + I = \pi \int_0^\pi \frac{\sin x}{1 + \sin x} dx$$

$$2I = \pi \int_0^\pi \frac{\sin x (1 - \sin x)}{(1 + \sin x)(1 - \sin x)} dx$$

$$2I = \pi \int_0^\pi \frac{\sin x - \sin^2 x}{(1 - \sin^2 x)} dx$$

$$2I = \pi \int_0^\pi \frac{\sin x - \sin^2 x}{(\cos^2 x)} dx$$

$$2I = \pi \int_0^\pi \left[\frac{\sin x}{\cos x} \cdot \frac{1}{\cos x} - \frac{\sin^2 x}{\cos^2 x} \right] dx$$

$$2I = \pi \int_0^\pi [\tan x \cdot \sec x - \tan^2 x] dx$$

$$2I = \pi \int_0^\pi \tan x \cdot \sec x dx - \pi \int_0^\pi \tan^2 x dx$$

$$\int \tan x \cdot \sec x dx = \sec x + c$$

$$2I = \pi [\sec x]_0^\pi - \pi \int_0^\pi (\sec^2 x - 1) dx$$

$$\int \sec^2 x dx = \tan x + c$$

$$2I = \pi [\sec \pi - \sec 0] - \pi [\tan x]_0^\pi - \pi [x]_0^\pi$$

$$2I = \pi[-1 - 1] - \pi[0 - 0] + \pi[\pi - 0]$$

$$2I = -2\pi + \pi^2$$

$$I = \frac{\pi^2}{2} - \pi$$

Laq Q.No 23

4. $\int_0^\pi \frac{x}{1 + \sin x} dx$

Sol:

$$I = \int_0^\pi \frac{x}{1 + \sin x} dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^\pi \frac{(\pi-x)}{1 + \sin(\pi-x)} dx$$

$$I = \int_0^\pi \frac{(\pi-x)}{1 + \sin x} dx$$

$$I = \int_0^\pi \frac{\pi}{1 + \sin x} dx - \int_0^\pi \frac{x}{1 + \sin x} dx$$

$$I + I = \pi \int_0^\pi \frac{1}{1 + \sin x} dx$$

$$2I = \pi \int_0^\pi \frac{(1 - \sin x)}{(1 + \sin x)(1 - \sin x)} dx$$

$$2I = \pi \int_0^\pi \frac{1 - \sin x}{(1 - \sin^2 x)} dx$$

$$2I = \pi \int_0^\pi \frac{1 - \sin x}{(\cos^2 x)} dx$$

$$2I = \pi \int_0^\pi \left[\frac{1}{\cos^2 x} - \frac{\sin x}{\cos x} \cdot \frac{1}{\cos x} \right] dx$$

$$2I = \pi \int_0^\pi [\sec^2 x - \tan x \cdot \sec x] dx$$

$$\int \tan x \cdot \sec x dx = \sec x + c$$

$$\int \sec^2 x dx = \tan x + c$$

$$2I = \pi [\tan x]_0^\pi - \pi [\sec x]_0^\pi$$

$$2I = \pi[\tan \pi - \tan 0] - \pi[\sec \pi - \sec 0]$$

$$2I = \pi[0 - 0] - \pi[-1 - 1]$$

$$2I = 2\pi$$

$$I = \pi$$

Definite integrals

5. $\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx$

Sol:

$$I = \int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^\pi \frac{(\pi-x) \sin(\pi-x)}{1 + \cos^2(\pi-x)} dx$$

$$I = \int_0^\pi \frac{(\pi-x) \sin x}{1 + \cos^2 x} dx$$

$$I = \int_0^\pi \frac{\pi \sin x}{1 + \cos^2 x} dx - \int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx$$

$$I + I = \pi \int_0^\pi \frac{\sin x}{1 + \cos^2 x} dx$$

$$2I = \pi \int_0^\pi \frac{\sin x}{1 + \cos^2 x} dx$$

let $\cos x = t \Rightarrow -\sin x dx = dt$

or

$$\sin x dx = -dt$$

$$L.L: x=0 \Rightarrow t=1; \quad U.L: x=\pi \Rightarrow t=-1$$

$$\Rightarrow 2I = \pi \int_1^{-1} \frac{-1}{1+t^2} dt$$

$$\Rightarrow 2I = \pi \int_{-1}^1 \frac{1}{1+t^2} dt$$

$$2I = \pi [\tan^{-1} t]$$

$$2I = \pi [\tan^{-1}(1) - \tan^{-1}(-1)]$$

$$2I = \pi \left[\frac{\pi}{4} + \frac{\pi}{4} \right]$$

$$2I = \pi \left[2 \cdot \frac{\pi}{4} \right] = \pi \left[\frac{\pi}{2} \right]$$

$$I = \frac{\pi^2}{4}$$

Laq Q.No 23

6. $\int_0^\pi \frac{x \sin^3 x}{1 + \cos^2 x} dx$

Sol:

$$I = \int_0^\pi \frac{x \sin^3 x}{1 + \cos^2 x} dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^\pi \frac{(\pi-x) \sin^3(\pi-x)}{1 + \cos^2(\pi-x)} dx$$

$$I = \int_0^\pi \frac{(\pi-x) \sin^3 x}{1 + \cos^2 x} dx$$

$$I = \int_0^\pi \frac{\pi \sin^3 x}{1 + \cos^2 x} dx - \int_0^\pi \frac{x \sin^3 x}{1 + \cos^2 x} dx$$

$$I + I = \pi \int_0^\pi \frac{\sin^3 x}{1 + \cos^2 x} dx = \pi \int_0^\pi \frac{\sin^3 x}{1 + \cos^2 x} dx$$

$$\text{let } \cos x = t \Rightarrow -\sin x dx = dt$$

or

$$\sin x dx = -dt$$

$$\sin^2 x = 1 - \cos^2 x = 1 - t^2$$

$$\sin^3 x = -(1 - t^2) dt$$

$$L.L: x=0 \Rightarrow t=1; \quad U.L: x=\pi \Rightarrow t=-1$$

$$\Rightarrow 2I = \pi \int_1^{-1} \frac{-(1-t^2)}{1+t^2} dt$$

$$\Rightarrow 2I = \pi \int_1^{-1} \frac{t^2-1}{1+t^2} dt = \pi \int_1^{-1} \frac{t^2+1-2}{1+t^2} dt$$

$$\Rightarrow 2I = \pi \int_1^{-1} \left[1 - \frac{2}{1+t^2} \right] dt$$

$$2I = \pi [t - 2\tan^{-1} t]_1^{-1}$$

$$2I = \pi [-1 - 1] - 2\pi [\tan^{-1}(-1) - \tan^{-1}(1)]$$

$$2I = -2\pi + 2\pi \left[\frac{\pi}{4} + \frac{\pi}{4} \right]$$

$$2I = -2\pi + 2\pi \left[\frac{\pi}{2} \right]$$

$$I = -\pi + \frac{\pi^2}{2} = \frac{\pi}{2}(\pi - 2)$$

Definite integrals

$$7. \int_0^{\pi/2} \frac{x}{\cos x + \sin x} dx$$

sol:

$$I = \int_0^{\pi/2} \frac{x}{\cos x + \sin x} dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^{\pi/2} \frac{(\frac{\pi}{2}-x)}{\cos(\frac{\pi}{2}-x)+\sin(\frac{\pi}{2}-x)} dx$$

$$I = \int_0^{\pi/2} \frac{(\frac{\pi}{2}-x)}{\sin x + \cos x} dx$$

$$= \int_0^{\pi/2} \frac{\frac{\pi}{2}}{\cos x + \sin x} dx - \int_0^{\pi/2} \frac{x}{\cos x + \sin x} dx$$

$$I + I = \frac{\pi}{2} \int_0^{\pi/2} \frac{1}{\cos x + \sin x} dx$$

$$2I = \frac{\pi}{2} \int_0^{\pi/2} \frac{1}{\cos x + \sin x} dx$$

Let $t = \tan\left(\frac{x}{2}\right)$;
 $dx = \frac{2dt}{1+t^2}$;
 $\cos x = \frac{1-t^2}{1+t^2}$
 $\sin x = \frac{2t}{1+t^2}$

$$2I = \frac{\pi}{2} \int_0^1 \frac{1}{\left(\frac{1-t^2}{1+t^2}\right) + \left(\frac{2t}{1+t^2}\right)} \left(\frac{2dt}{1+t^2}\right)$$

$$2I = \frac{\pi}{2} \int_0^1 \frac{1}{\left[\frac{1-t^2+2t}{1+t^2}\right]} \left(\frac{2dt}{1+t^2}\right)$$

$$2I = \frac{\pi}{2} 2 \int_0^1 \frac{1}{-(t^2-2t-1)} dt$$

$$I = \frac{\pi}{2} \int_0^1 \frac{1}{-[t^2-2t+(1)^2-(1)^2-1]} dt$$

Laq Q.No 23

$$I = \frac{\pi}{2} \int_0^1 \frac{1}{-(t-1)^2 - (\sqrt{2})^2} dt$$

$$I = \frac{\pi}{2} \int_0^1 \frac{1}{[(\sqrt{2})^2 - (t-1)^2]} dt$$

$$\therefore \int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + c$$

$$2I = \frac{\pi}{2} \frac{1}{2(\sqrt{2})} \left[\log \left| \frac{\sqrt{2}+t-1}{\sqrt{2}-t+1} \right| \Big|_0^1 \right]$$

$$I = \frac{\pi}{2} \frac{1}{2\sqrt{2}} \left[\log \left| \frac{\sqrt{2}+1-1}{\sqrt{2}-1+1} \right| - \log \left| \frac{\sqrt{2}-1}{\sqrt{2}+1} \right| \right]$$

$$I = \frac{\pi}{2} \frac{1}{2\sqrt{2}} \left[\log |1| + \log \left| \frac{\sqrt{2}+1}{\sqrt{2}-1} \right| \right]$$

$$I = \frac{\pi}{2} \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2}+1}{\sqrt{2}-1} \right|$$

$$I = \frac{\pi}{2} \frac{1}{2\sqrt{2}} \log \left| \frac{(\sqrt{2}+1)(\sqrt{2}+1)}{(\sqrt{2}-1)(\sqrt{2}+1)} \right|$$

$$I = \frac{\pi}{2} \frac{1}{2\sqrt{2}} \log \frac{(\sqrt{2}+1)^2}{\sqrt{2}^2 - 1^2}$$

$$I = \frac{2}{2\sqrt{2}} \log \frac{(\sqrt{2}+1)}{2-1}$$

$$I = \frac{\pi}{2\sqrt{2}} \log(\sqrt{2} + 1)$$

Definite integrals

$$8. \int_0^{\pi/2} \frac{\sin^2 x}{\cos x + \sin x} dx$$

sol:

$$I = \int_0^{\pi/2} \frac{\sin^2 x}{\cos x + \sin x} dx \dots (1)$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^{\pi/2} \frac{\sin^2(\frac{\pi}{2}-x)}{\cos(\frac{\pi}{2}-x)+\sin(\frac{\pi}{2}-x)} dx$$

$$I = \int_0^{\pi/2} \frac{\cos^2 x}{\sin x + \cos x} dx \dots (2)$$

Adding (1) & (2)

$$= \int_0^{\pi/2} \frac{\sin^2 x}{\cos x + \sin x} dx + \int_0^{\pi/2} \frac{\cos^2 x}{\cos x + \sin x} dx$$

$$I + I = \int_0^{\pi/2} \frac{\sin^2 x + \cos^2 x}{\cos x + \sin x} dx$$

$$2I = \int_0^{\pi/2} \frac{1}{\cos x + \sin x} dx$$

$$\text{Let } t = \tan\left(\frac{x}{2}\right);$$

$$dx = \frac{2dt}{1+t^2};$$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$\sin x = \frac{2t}{1+t^2}$$

$$I = \int_0^1 \frac{1}{[(t-1)^2 - (\sqrt{2})^2]} dt$$

$$I = \int_0^1 \frac{1}{[(\sqrt{2})^2 - (t-1)^2]} dt$$

$$\therefore \int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + c$$

$$2I = \frac{1}{2(\sqrt{2})} \left[\log \left| \frac{\sqrt{2}+t-1}{\sqrt{2}-t+1} \right| \Big|_0^1 \right]$$

$$I = \frac{1}{2\sqrt{2}} \left[\log \left| \frac{\sqrt{2}+1-1}{\sqrt{2}-1+1} \right| - \log \left| \frac{\sqrt{2}-1}{\sqrt{2}+1} \right| \right]$$

$$I = \frac{1}{2\sqrt{2}} \left[\log |1| + \log \left| \frac{\sqrt{2}+1}{\sqrt{2}-1} \right| \right]$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2}+1}{\sqrt{2}-1} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{(\sqrt{2}+1)(\sqrt{2}+1)}{(\sqrt{2}-1)(\sqrt{2}+1)} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \frac{(\sqrt{2}+1)^2}{\sqrt{2}^2 - 1^2}$$

$$I = \frac{2}{2\sqrt{2}} \log \frac{(\sqrt{2}+1)}{2-1}$$

$$I = \frac{1}{\sqrt{2}} \log(\sqrt{2} + 1)$$

$$2I = \int_0^1 \frac{1}{\left(\frac{1-t^2}{1+t^2} \right) + \left(\frac{2t}{1+t^2} \right)} \left(\frac{2dt}{1+t^2} \right)$$

$$2I = \int_0^1 \frac{1}{\left[\frac{1-t^2+2t}{1+t^2} \right]} \left(\frac{2dt}{1+t^2} \right)$$

$$2I = 2 \int_0^1 \frac{1}{-(t^2-2t-1)} dt$$

$$I = \int_0^1 \frac{1}{-[t^2-2t+(1)^2-(1)^2-1]} dt$$

Laq Q.No 23

Definite integrals

9. $\int_3^7 \sqrt{\frac{7-x}{x-3}} dx.$

Sol:

$$\begin{aligned} let \ x &= 3\cos^2\theta + 7\sin^2\theta \\ dx &= 8\sin\theta\cos\theta.d\theta \end{aligned}$$

$$L.L: x = 3 \Rightarrow \theta = 0$$

$$U.L: x = 7 \Rightarrow \theta = \frac{\pi}{2}$$

$$\begin{aligned} 7 - x &= 7 - 3\cos^2\theta - 7\sin^2\theta \\ &= 7(1 - \sin^2\theta) - 3\cos^2\theta \\ &= 7(\cos^2\theta) - 3\cos^2\theta \\ &= 4\cos^2\theta \end{aligned}$$

$$\begin{aligned} x - 3 &= 3\cos^2\theta + 7\sin^2\theta - 3 \\ &= 7\sin^2\theta - 3(1 - \cos^2\theta) \\ &= 7\sin^2\theta - 3(\sin^2\theta) \\ &= 4\sin^2\theta \end{aligned}$$

$$\begin{aligned} \int_3^7 \sqrt{\frac{7-x}{x-3}} dx &= \int_0^{\pi/2} \sqrt{\frac{4\cos^2\theta}{4\sin^2\theta}} 8\sin\theta\cos\theta d\theta \\ &= 8 \int_0^{\pi/2} \cos^2\theta d\theta \end{aligned}$$

$$\int_0^{\pi/2} \cos^n x dx = \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \cdots \frac{\pi}{2}$$

$$= 8 \cdot \left(\frac{1}{2}\right) \frac{\pi}{2} = 2\pi$$

10. $\int_4^9 \frac{1}{\sqrt{(9-x)(x-4)}} dx.$

Sol:

$$\begin{aligned} let \ x &= 4\cos^2\theta + 9\sin^2\theta \\ dx &= 10\sin\theta\cos\theta.d\theta \end{aligned}$$

$$L.L: x = 4 \Rightarrow \theta = 0$$

$$U.L: x = 9 \Rightarrow \theta = \frac{\pi}{2}$$

$$\begin{aligned} 9 - x &= 9 - 4\cos^2\theta - 9\sin^2\theta \\ &= 9(1 - \sin^2\theta) - 3\cos^2\theta \\ &= 9(\cos^2\theta) - 4\cos^2\theta \\ &= 5\cos^2\theta \end{aligned}$$

$$x - 4 = 4\cos^2\theta + 9\sin^2\theta - 4$$

$$\begin{aligned} &= 9\sin^2\theta - 4(1 - \cos^2\theta) \\ &= 9\sin^2\theta - 4(\sin^2\theta) \\ &= 5\sin^2\theta \end{aligned}$$

$$\begin{aligned} \int_4^9 \frac{1}{\sqrt{(9-x)(x-4)}} dx &= \int_0^{\pi/2} \frac{1}{\sqrt{5\cos^2\theta 5\sin^2\theta}} 10\sin\theta\cos\theta d\theta \\ &= \int_0^{\pi/2} \frac{1}{5\sin\theta\cos\theta} 10\sin\theta\cos\theta d\theta \end{aligned}$$

$$I = 2 \int_0^{\pi/2} 1 dx$$

$$= 2[x]_0^{\pi/2} = \frac{2\pi}{2} = \pi$$

11. Evaluate $\int_a^b \sqrt{(x-a)(x-b)} dx$

$$\text{Sol: } \int_a^b \sqrt{-(x-a)(x-b)} dx$$

$$I = \int_a^b \sqrt{-[x^2 - (a+b)x + ab]} dx$$

$$= \int_a^b \sqrt{-\left[x^2 - (a+b)x + \frac{[a+b]^2}{4} - \frac{[a+b]^2}{4} + ab\right]} dx$$

$$= \int_a^b \sqrt{-\left[\left[x - \frac{a+b}{2}\right]^2 - \frac{[b-a]^2}{4}\right]} dx$$

$$= \int_a^b \sqrt{\left[\left[\frac{b-a}{2}\right]^2 - \left[x - \frac{a+b}{2}\right]^2\right]} dx$$

$$\therefore \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + C$$

$$I = \left[\frac{x - \frac{a+b}{2}}{2} \sqrt{(a-x)(b-x)} + \frac{[b-a]^2}{4 \cdot 2} \sin^{-1} \left(\frac{x - \frac{a+b}{2}}{\frac{b-a}{2}} \right) \right]_a^b$$

$$= \left[\frac{b - \frac{a+b}{2}}{2} \sqrt{(a-b)(b-b)} + \frac{[b-a]^2}{4 \cdot 2} \sin^{-1} \left(\frac{b - \frac{a+b}{2}}{\frac{b-a}{2}} \right) \right]$$

$$- \left[\frac{a - \frac{a+b}{2}}{2} \sqrt{(a-a)(b-a)} + \frac{[b-a]^2}{4 \cdot 2} \sin^{-1} \left(\frac{a - \frac{a+b}{2}}{\frac{b-a}{2}} \right) \right]$$

$$= \left[\frac{[b-a]^2}{8} \sin^{-1} \left(\frac{b-a}{\frac{b-a}{2}} \right) \right] - \left[\frac{[b-a]^2}{8} \sin^{-1} \left(\frac{a-b}{\frac{b-a}{2}} \right) \right]$$

$$= \left[\frac{[b-a]^2}{8} \sin^{-1} 1 \right] - \left[\frac{[b-a]^2}{8} \sin^{-1} (-1) \right]$$

$$= \left[\frac{[b-a]^2}{8} \left(\frac{\pi}{2} \right) \right] + \left[\frac{[b-a]^2}{8} \left(\frac{\pi}{2} \right) \right] = \frac{\pi [b-a]^2}{8}$$

Laq Q.No 23

12. $\int_0^\pi x \cdot \sin^7 x \cos^6 x dx$

Sol:

$$I = \int_0^\pi x \cdot \sin^7 x \cos^6 x dx$$

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^\pi (\pi - x) \sin^7(\pi - x) \cos^6(\pi - x) dx$$

$$I = \int_0^\pi (\pi - x) \sin^7 x \cos^6 x dx$$

$$I = \int_0^\pi \pi \sin^7 x \cos^6 x dx - \int_0^\pi x \sin^7 x \cos^6 x dx$$

$$I = \int_0^\pi \pi \sin^7 x \cos^6 x dx - I$$

$$I + I = \pi \int_0^\pi \sin^7 x \cos^6 x dx$$

$$\therefore \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx$$

$$2I = 2\pi \int_0^{\pi/2} \sin^7 x \cos^6 x dx$$

$$I = \pi \int_0^{\pi/2} \sin^7 x \cos^6 x dx$$

$$\int_0^{\pi/2} \cos^n x \sin^m x dx = \frac{(n-1)(n-3)(n-5) \dots (m-1)(m-3) \dots}{(m+n)(m+n-2)(m+n-4) \dots}$$

$$I = \pi \frac{6.4.2.5.3.1}{13.11.9.7.5.3.1}$$

$$I = \pi \frac{6.4.2}{13.11.9.7}$$

$$I = \frac{16\pi}{3003}$$

Definite integrals

13. Find the area enclosed by the curves

$$y^2 = 4ax \text{ and } x^2 = 4by.$$

Sol:

Given eq'n

$$y^2 = 4ax \Rightarrow y = \sqrt{4ax} \dots (1)$$

$$x^2 = 4by \Rightarrow y = \frac{x^2}{4b} \dots (2)$$

solving (1)and (2)

$$\sqrt{4ax} = \frac{x^2}{4b} \text{ S.O.B}$$

$$\Rightarrow 4ax = \frac{x^4}{16b^2}$$

$$\Rightarrow 64ab^2x = x^4 \Rightarrow 64ab^2x - x^4 = 0$$

$$\Rightarrow x(64ab^2 - x^3) = 0$$

$$x = 0 \text{ or } x^3 = 64ab^2$$

$$\Rightarrow x = 4a^{1/3}b^{2/3} = u$$

$$\text{Required Area} = \int_0^u [(1) - (2)] dx$$

$$= \int_0^u \left[\sqrt{4ax} - \frac{x^2}{4b} \right] dx$$

$$= \int_0^u \left[2\sqrt{ax^{1/2}} - \frac{x^2}{4b} \right] dx$$


$$= \left[\frac{2\sqrt{ax^{3/2}}}{3/2} - \frac{x^3}{4b3} \right]_0^u$$

$$= \frac{4}{3}\sqrt{a}[u^{3/2} - 0^2] - \frac{1}{12}[u^3 - 0^3]$$

$$= \frac{4}{3}\sqrt{a} \left[\left[4a^{\frac{1}{3}}b^{\frac{2}{3}} \right]^{\frac{3}{2}} - 0^2 \right] - \frac{1}{12} \left[[4a^{1/3}b^{2/3}]^3 - 0^3 \right]$$

$$= \frac{4}{3}[8ab] - \frac{1}{12b}[64ab^2]$$

$$= \frac{32ab}{3} - \frac{16ab}{3} = \frac{16ab}{3} \text{ sq. units}$$



14. Find the area enclosed by the curves

$$y = 4 - 2x \text{ and } y = x^2 - 5x.$$

Sol:

Given eq'n

$$y = 4 - 2x \dots (1)$$


$$y = x^2 - 5x \dots (1)$$

solving (1)and (2)

$$x^2 - 5x = 4 - 2x$$

$$\Rightarrow x^2 - 3x - 4 = 0$$

$$\Rightarrow x^2 + 1x - 4x - 4 = 0$$



$$\Rightarrow x(x+1) - 4(x+1) = 0$$

$$x = 4 \text{ or } x = -1$$

$$\text{Required Area} = \int_{-1}^4 [(1) - (2)] dx$$

$$= \int_{-1}^4 [4 - 2x - x^2 + 5x] dx$$

$$= \int_{-1}^4 [4 + 3x - x^2] dx$$

$$= \left[4x + \frac{3x^2}{2} - \frac{x^3}{3} \right]_{-1}^4$$

$$= 4[4 + 1] + \frac{3}{2}[4^2 - (-1)^2] - \frac{1}{3}[4^3 - (-1)^3]$$

$$= 20 + \frac{3}{2}(16 - 1) - \frac{1}{3}(64 + 1)$$

$$= 20 + \frac{45}{2} - \frac{65}{3}$$

$$= \frac{120 + 135 - 135}{6}$$

$$= \frac{125}{6} \text{ sq. units}$$

15. Find the area enclosed by the curves

$$y^2 = 4x \text{ and } y^2 = 4(4 - x).$$

Sol:

Given eq'n

$$y^2 = 4x \Rightarrow y = \sqrt{4x} \dots (1)$$

$$y^2 = 4(4 - x) \Rightarrow y = \sqrt{4(4 - x)} \dots (2)$$

solving (1)and (2)

$$4x = 4(4 - x)$$

$$\Rightarrow x = 4 - x$$

$$\Rightarrow 2x = 4$$

$$x = 2$$

Sub x=2 in (1)

$$y^2 = 4x = 4(2) = 8$$

$$y = \sqrt{8} = \pm 2\sqrt{2}$$

Two parabolas are symmetric about X-axis

$$\text{Required Area} = 2 \left[\int_0^2 (1) dx + \int_2^4 (2) dx \right]$$

$$= 2 \left[\int_0^2 \sqrt{4x} dx + \int_2^4 \sqrt{4(4-x)} dx \right]$$

$$= 2 \left[2 \int_0^2 x^{1/2} dx + 2 \int_2^4 (4-x)^{1/2} dx \right]$$


$$= 4 \left[\frac{x^{3/2}}{3/2} \right]_0^2 + 4 \left[\frac{(4-x)^{3/2}}{-3/2} \right]_2^4$$

$$= \frac{8}{3} [2^{3/2} - 0^{3/2}] - \frac{8}{3} [(4-4)^{3/2} - (4-2)^{3/2}]$$

$$= \frac{8}{3} [2\sqrt{2}] + \frac{8}{3} [2\sqrt{2}]$$

$$= \frac{16\sqrt{2}}{2} + \frac{16\sqrt{2}}{2}$$

$$= \frac{32\sqrt{2}}{2} \text{ sq. units}$$



16. Find the area enclosed by the curves

$$y = 2 - x^2 \text{ and } y = x^2.$$

Sol:

Given eq'n

$$y = 2 - x^2 \dots (1)$$

$$y = x^2 \dots (2)$$

solving (1)and (2)


$$2 - x^2 = x^2$$

$$2 = 2x^2$$

$$x^2 = 1$$

$$x = \pm 1$$

$$\therefore x = 1 \text{ or } x = -1$$



$$\text{Required Area} = \int_{-1}^1 [(1) - (2)] dx$$

$$= \int_{-1}^1 [2 - x^2 - x^2] dx$$

$$= \int_{-1}^1 [2 - 2x^2] dx$$

$$= \left[2x - 2 \frac{x^3}{3} \right]_{-1}^1$$

$$= 2[1 + 1] - \frac{2}{3}[(1)^3 - (-1)^3]$$

$$= 4 - \frac{2}{3}(1 + 1)$$

$$= 4 - \frac{4}{3} = \frac{12-4}{3}$$

$$= \frac{8}{3} \text{ sq. unit}$$

17. Show that the area of the region bounded by

$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is πab . also deduce the area of the circle $x^2 + y^2 = a^2$.

Sol: Given eq'n of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \dots \dots \dots (1)$$

$$\Rightarrow \frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$\Rightarrow \frac{y^2}{b^2} = \frac{1}{a^2} [a^2 - x^2]$$

$$\Rightarrow y^2 = \frac{b^2}{a^2} [a^2 - x^2]$$

$$\Rightarrow y = \pm \frac{b}{a} \sqrt{a^2 - x^2}$$

Ellipse is symmetric about both the axes.

Required area = 4 area of shaded region

$$\text{Area} = \int_0^a y dx$$

$$= 4 \int_0^{\frac{a}{a}} \frac{b}{a} \sqrt{a^2 - x^2} dx$$

$$= 4 \frac{b}{a} \int_0^a \sqrt{a^2 - x^2} dx$$

$$\therefore \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + c$$

$$= \frac{4b}{a} \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) \right]_0^a$$

$$= \frac{4b}{a} \left[\frac{a}{2} \sqrt{a^2 - a^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{a}{a} \right) - 0 - 0 \right]$$

$$= \frac{4b}{a} \frac{a^2}{2} \sin^{-1} (1)$$

$$= \frac{2ab\pi}{2}$$

$$= \pi ab \text{ sq units}$$

If $a=b$ the ellipse becomes a circle

\therefore Area of the circle $x^2 + y^2 = a^2$.

is $\pi a \cdot a = \pi a^2 \text{ sq . units}$

