PAIR OF LINES-SECOND DEGREE GENERAL EQUATION

THEOREM

If the equation $S = ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of straight lines then

i)
$$\Delta \equiv abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$
 and (ii) $h^2 \ge ab$, $g^2 \ge ac$, $f^2 \ge bc$

Proof:

Let the equation S = 0 represent the two lines $l_1x + m_1y + n_1 = 0$ and $l_2x + m_2y + n_2 = 0$.

Then

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c$$

$$\equiv (l_{1}x + m_{1}y + n_{1})(l_{2}x + m_{2}y + n_{2}) = 0$$

Equating the co-efficients of like terms, we get

$$l_1 l_2 = a$$
, $l_1 m_2 + l_2 m_1 = 2h$, $m_1 m_2 = b$, and $l_1 n_2 + l_2 n_1 = 2g$, $m_1 n_2 + m_2 n_1 = 2f$, $n_1 n_2 = c$

(i) Consider the product
$$(2h)(2g)(2f)$$

$$= (l_1m_2 + l_2m_1)(l_1n_2 + l_2n_1)(m_1n_2 + m_2n_1)$$

$$= l_1l_2\left(m_1^2n_2^2 + m_2^2n_1^2\right) + m_1m_2\left(l_1^2n_2^2 + l_2^2n_1^2\right) + n_1n_2\left(l_1^2m_2^2 + l_2^2m_1^2\right) + 2l_1l_2m_1m_2n_1n_2$$

$$= l_1l_2\left[(m_1n_2 + m_2n_1)^2 - 2m_1m_2n_1n_2\right] + m_1m_2\left[(l_1n_2 + l_2n_1)^2 - 2l_1l_2n_1n_2\right]$$

$$+ n_1n_2\left[(l_1m_2 + l_2m_1)^2 - 2l_1l_2m_1m_2\right] + 2l_1l_2m_1m_2n_1n_2$$

$$= a(4f^2 - 2bc) + b(4g^2 - 2ac) + c(4h^2 - 2ab)$$

$$8fgh = 4[af^2 + bg^2 + ch^2 - abc]$$

$$\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

ii)
$$h^2 - ab = \left(\frac{l_1 m_2 + l_2 m_1}{2}\right)^2 - l_1 l_2 m_1 m_2 = \frac{(l_1 m_2 + l_2 m_1)^2 - 4 - l_1 l_2 m_1 m_2}{4}$$

$$= \frac{(l_1 m_2 - l_2 m_1)^2}{4} \ge 0$$

Similarly we can prove $g^2 \ge ac$ and $f^2 \ge bc$

NOTE:

If $\Delta = abc + 2fgh - af^2 - bg^2 - ch^2 = 0$, $h^2 \ge ab$, $g^2 \ge ac$ and $f^2 \ge bc$, then the equation $S = ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of straight lines

CONDITIONS FOR PARALLEL LINES-DISTANCE BETWEEN THEM

THEOREM

If $S = ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of parallel lines then $h^2 = ab$ and $bg^2 = af^2$. Also the distance between the two parallel lines is

$$2\sqrt{\frac{g^2-ac}{a(a+b)}} \text{ (or) } 2\sqrt{\frac{f^2-bc}{b(a+b)}}$$

Proof:

Let the parallel lines represented by S = 0 be $lx + my + n_1 = 0$ -- (1) $lx + my + n_2 = 0$ -- (2)

$$\therefore ax^2 + 2hxy + 2gx + 2fy + c$$

$$\equiv (lx + my + n_1)(lx + my + n_2)$$

Equating the like terms

$$l^2 = a - (3)$$
 $2lm = 2h - (4)$

$$m^2 = b - (5)$$
 $l(n_1 + n_2) = 2g - (6)$

$$m(n_1 + n_2) = 2f - (7)$$
 $n_1n_2 = c - (8)$

From (3) and (5), $l^2m^2 = ab$ and from (4) $h^2 = ab$.

Dividing (6) and (7)
$$\frac{l}{m} = \frac{g}{f} \Rightarrow \frac{l^2}{m^2} = \frac{g^2}{f^2}$$
,

$$\therefore \frac{a}{b} = \frac{g^2}{f^2} \Rightarrow bg^2 = af^2$$

Distance between the parallel lines (1) and (2) is

$$= \left| \frac{n_1 - n_2}{\sqrt{(l^2 + m^2)}} \right| = \frac{\sqrt{(n_1 + n_2)^2 - 4n_1 n_2}}{\sqrt{l^2 + m^2}}$$

$$= \frac{\sqrt{(4g^2 / l^2) - 4c}}{\sqrt{a + b}} \text{ or } \frac{\sqrt{(4f^2 / m^2) - 4c}}{\sqrt{a + b}}$$

$$= 2\sqrt{\frac{g^2 - ac}{a(a + b)}} \text{ (or) } 2\sqrt{\frac{f^2 - bc}{b(a + b)}}$$

POINT OF INTERSECTION OF PAIR OF LINES THEOREM

The point of intersection of the pair of lines represented by

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$$
 when $h^{2} > ab$ is $\left(\frac{hf - bg}{ab - h^{2}}, \frac{gh - af}{ab - h^{2}}\right)$

Proof:

Let the point of intersection of the given pair of lines be (x_1, y_1) . Transfer the origin to (x_1, y_1) without changing the direction of the axes.

Let (X,Y) represent the new coordinates of

$$(x, y)$$
. Then $x = X + x_1$ and $y = Y + y_1$.

Now the given equation referred to new axes will be

$$a(X + x_1)^2 + 2h(X + x_1)(Y + y_1) + b(Y + y_1)^2 + 2g(X + x_1) + 2f(Y + y_1) + c = 0$$

$$\Rightarrow aX^2 + 2hXY + bY^2 + 2X(ax_1 + hy_1 + g) + 2Y(hx_1 + by_1 + f)$$

$$+(ax_1^2 + 2hx_1y_1 + by_1^2 + 2gx_1 + 2fy_1 + c) = 0$$

Since this equation represents a pair of lines passing through the origin it should be a homogeneous second degree equation in X and Y. Hence the first degree terms and the constant term must be zero. Therefore,

$$ax_1 + hy_1 + g = 0$$
 -- (1)

$$hx_1 + by_1 + f = 0$$
 -- (2)

$$ax_1^2 + 2hx_1y_1 + by_1^2 + 2gx_1 + 2fy_1 + c = 0$$
 -- (3)

But (3) can be rearranged as

$$x_1(ax_1 + hy_1 + g) + y_1(hx_1 + by_1 + f) + (gx_1 + fy_1 + c) = 0$$

$$\Rightarrow gx_1 + fy_1 + c = 0 - -(4)$$

Solving (1) and (2) for x_1 and y_1

$$\frac{x_1}{hf - bg} = \frac{y}{gh - af} = \frac{1}{ab - h^2}$$

$$\therefore x_1 = \frac{hf - bg}{ab - h^2} \text{ and } y_1 = \frac{gh - af}{ab - h^2}$$

Hence the point of intersection of the given pair of lines is $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$

THEOREM


If the pair of lines $ax^2 + 2hxy + by^2 = 0$ and the pair of lines

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$
 form a rhombus then $(a-b)fg + h(f^2 - g^2) = 0$.

Proof:

The pair of lines $ax^2 + 2hxy + by^2 = 0$ -- (1) is parallel to the lines

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$
 -- (2)



Now the equation

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c + \lambda(ax^{2} + 2hxy + by^{2}) = 0$$

Represents a curve passing through the points of intersection of (1) and (2).

Substituting $\lambda = -1$, in (3) we obtain 2gx + 2fy + c = 0 ...(4) Equation (4) is a straight line passing through A and B and it is the diagonal \overrightarrow{AB}

The point of intersection of (2) is $C = \left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$

$$\Rightarrow$$
 Slope of $\overrightarrow{OC} = \frac{gh - af}{hf - bg}$

In a rhombus the diagonals are perpendicular \Rightarrow (Slope of \overrightarrow{OC})(Slope of \overrightarrow{AB}) = -1

$$\Rightarrow \left(\frac{gh - af}{hf - bg}\right) \left(-\frac{g}{f}\right) = -1$$

$$\Rightarrow g^2h - afg = hf^2 - bfg$$

$$\Rightarrow (a-b)fg + h(f^2 - g^2) = 0$$


$$\frac{g^2 - f^2}{a - b} = \frac{fg}{h}$$

THEOREM

If $ax^2 + 2hxy + by^2 = 0$ be two sides of a parallelogram and px + qy = 1 is one diagonal, then the other diagonal is y(bp - hq) = x(aq - hp)

proof:

Let $P(x_1, y_1)$ and $Q(x_2, y_2)$ be the points where the digonal



px + qy = 1 meets the pair of lines.

 \overline{OR} and \overline{PQ} biset each other at $M(\alpha, \beta)$.

$$\therefore \alpha = \frac{x_1 + x_2}{2} \text{ and } \beta = \frac{y_1 + y_2}{2}$$

Eliminating y from $ax^2 + 2hxy + by^2 = 0$ -- (1)

and
$$px + qy = 1 \qquad --(2)$$
$$ax^{2} + 2hx \left(\frac{1-px}{q}\right) + b\left(\frac{1-px}{q}\right)^{2} = 0$$

$$\Rightarrow x^2(aq^2 - 2hpq + bp^2) + 2x(hp - bp) + b = 0$$

The roots of this quadratic equation are x₁ and x₂ where

$$x_1 + x_2 = -\frac{2(hq - bp)}{aq^2 - 2hpq - bp^2}$$

$$\Rightarrow \alpha = \frac{(bp - hq)}{(aq^2 - 2hpq + bp^2)}$$

Similarly by eliminating x from (1) and (2) a quadratic equation in y is obtained and y₁,

y₂ are its roots where

$$y_1 + y_2 = -\frac{2(hp - aq)}{aq^2 - 2hpq - np^2} \Rightarrow \beta = \frac{(aq - hp)}{(aq^2 - 2hpq + bp^2)}$$

Now the equation to the join of O(0, 0) and $M(\alpha, \beta)$ is $(y-0)(0-\alpha) = (x-0)(0-\beta)$

$$\Rightarrow \alpha y = \beta x$$

Substituting the values of α and β , the equation of the diagonal OR

is
$$y(bp-hq) = x(aq-hp)$$
.

EXERCISE 4B

I

1. Find the angle between the lines represented by $2x^2 + xy - 6y^2 + 7y - 2 = 0$.

Sol. Given equation is

$$2x^{2} + xy - 6y^{2} + 7y - 2 = 0$$
 Comparing with

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$$
 then

$$a = 2, b = -6, c = -2, g = 0, f = \frac{7}{2}, h = \frac{1}{2}$$

Angle between the lines is given by

$$\cos \alpha = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|2-6|}{\sqrt{(2+6)^2 + 1}} = \frac{4}{\sqrt{65}} \implies \alpha = \cos^{-1}\left(\frac{4}{\sqrt{65}}\right)$$

- 2. Prove that the equation $2x^2 + 3xy 2y^2 + 3x + y + 1 = 0$ represents a pair of perpendicular lines.
- **Sol.** From given equation a = 2, b = -2 and a + b = 2 + (-2) = 0
 - \Rightarrow angle between the lines is 90°. \therefore The given lines are perpendicular.

- 1. Prove that the equation $3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$ represents a pair of straight lines and find the co-ordinates of the point of intersection.
- **Sol.** The given equation is $3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$

Comparing with $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, we get

$$a = 3$$
 , $b = 2$, $c = 2$, $2f = 3$ $\Rightarrow f = \frac{5}{2}$

$$2g = 5 \implies g = \frac{5}{2},$$
 $2h = 7 \implies h = \frac{7}{2}$

$$\Delta = abc + 2fgh - af^2 - bg^2 - ch^2$$

$$=3(2)(2)+2.\frac{5}{2}.\frac{5}{2}.\frac{7}{2}-3.\frac{25}{4}-2.\frac{25}{4}-2.\frac{49}{4}$$

$$= \frac{1}{4} (48 + 175 - 75 - 50 - 98)$$

$$=\frac{1}{2}(223-223)=0$$

$$h^2 - ab = \left(\frac{7}{2}\right)^2 - 3.2 = \frac{49}{4} - 6 = \frac{25}{4} > 0$$

$$f^2 - bc = \left(\frac{5}{2}\right)^2 - 2.2 = \frac{25}{4} - 4 = \frac{9}{4} > 0$$

$$g^2 - ac = \left(\frac{5}{2}\right)^2 - 3.2 = \frac{25}{4} - 6 = \frac{1}{4} > 0$$

:. The given equation represents a pair of lines.

The point of intersection of the lines is $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$

$$= \left(\frac{\frac{7}{2} \cdot \frac{5}{2} - 2\frac{5}{2}}{6 - \frac{49}{4}}, \frac{\frac{5}{2} \cdot \frac{7}{2} - 3 \cdot \frac{5}{2}}{6 - \frac{49}{4}}\right) = \left(\frac{35 - 20}{24 - 29}, \frac{35 - 30}{24 - 49}\right)$$

$$=\left(\frac{+15}{-25}, \frac{5}{-28}\right) = \left(\frac{-3}{5}, -\frac{1}{5}\right)$$

Point of intersection is $p\left(\frac{-3}{5}, \frac{-1}{5}\right)$

- 2. Find the value of k, if the equation $2x^2 + kxy 6y^2 + 3x + y + 1 = 0$ represents a pair of straight lines. Find the point of intersection of the lines and the angle between the straight lines for this value of k.
- **Sol.** The given equation is $2x^2 + kxy 6y^2 + 3x + y + 1 = 0$

$$a = 2$$
, $b = -6$, $c = 1$, $f = \frac{1}{2}$, $2g = 3g = \frac{3}{2}$, $h = \frac{k}{2}$

Since the given equation is representing a pair of straight lines, therefore

$$\Delta = abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow -12 + 2 \cdot \frac{1}{2} \cdot \frac{3}{2} \cdot \left(+\frac{k}{2} \right) - 2 \cdot \frac{1}{4} + 6 \cdot \frac{9}{4} - \frac{k^2}{4} = 0$$

$$\Rightarrow -48 + 3k - 2 + 54 - k^2 = 0$$

$$\Rightarrow$$
 $-k^2 + 3k + 4 = 0 \Rightarrow k^2 - 3k - 4 = 0$

$$\Rightarrow (k-4)(k+1) = 0$$

$$\Rightarrow$$
 k = 4 or -1

Case (i) k = -1

Point of intersection is $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$

$$\left(\frac{+\frac{1}{2}\cdot\frac{1}{2}+6\cdot\frac{3}{2}}{-12-\frac{1}{4}},\frac{\frac{3}{2}\left(-\frac{1}{2}\right)-2\cdot\frac{1}{2}}{-12-\frac{1}{4}}\right) = \left(\frac{-1+36}{-49},\frac{-3-4}{-49}\right)$$

$$=\left(\frac{35}{-49}, \frac{-7}{-49}\right) = \left(\frac{-5}{7}, \frac{1}{7}\right)$$

Point of intersection is $\left(\frac{-5}{7}, \frac{1}{7}\right)$

Angle between the lines
$$= \cos \alpha = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|2-6|}{\sqrt{(2+6)^2 + 4}} = \left(\frac{4}{\sqrt{65}}\right)$$

Case (ii) k = 4

$$\left(\frac{2.\frac{1}{2} + 6.\frac{3}{2}}{-12 - 4}, \frac{\frac{3}{2}.2 - 2.\frac{1}{2}}{-12 - 4}\right) = \left(-\frac{5}{8}, -\frac{1}{8}\right)$$

Point of intersection is $P\left(-\frac{5}{8}, -\frac{1}{8}\right)$ and angle between the lines is

$$\cos \alpha = \frac{\left|a+b\right|}{\sqrt{\left(a-b\right)^2 + 4h^2}}$$

$$=\frac{|2-6|}{\sqrt{(2+6)^2+16}}=\frac{4}{4\sqrt{5}}=\frac{1}{\sqrt{5}}$$

$$\alpha = \cos^{-1}\left(\frac{1}{\sqrt{5}}\right)$$

- 3. Show that the equation $x^2 y^2 x + 3y 2 = 0$ represents a pair of perpendicular lines and find their equations.
- Sol. Given equation is $x^2 y^2 x + 3y 2 = 0 \Rightarrow a = 1, b = 1, c = -2 \text{ f} = \frac{3}{2}, g = -\frac{1}{2}$

$$h = 0$$

Now $\Delta = abc + 2fgh - af^2 - bg^2 - ch^2$

$$=1(-1)(-2)+0-1.\frac{9}{4}+1.\frac{1}{4}+0 = +2-\frac{9}{4}+\frac{1}{4}=0$$

$$h^2 - ab = 0 - 1(-1) = 1 > 0$$

$$f^2 - bc = \frac{9}{4} - 2 = \frac{1}{4} > 0$$

$$g^2 - ac = \frac{1}{4} + 2 = \frac{9}{4} > 0$$

And
$$a + b = 1 - 1 = 0$$

The given equation represent a pair of perpendicular lines.

Let
$$x^2 - y^2 - x + 3y - 2 = (x + y + c_1)(x - y + c_2)$$

Equating the coefficients of $x \Rightarrow c_1 + c_2 = -1$

Equating the co-efficient of $y \Rightarrow -c_1 + c_2 = 3$

Adding
$$2c_1 = 2 \Rightarrow c_2 = 1$$

$$c_1 + c_2 = -1 \Rightarrow c_1 + 1 = -1, c_1 = -2$$

Equations of the lines are x + y - 2 = 0 and x - y + 1 = 0

- 4. Show that the lines $x^2 + 2xy 35y^2 4x + 44y 12 = 0$ are 5x + 2y 8 = 0 are concurrent.
- **Sol.** Equation of the given lines are $x^2 + 2xy 35y^2 4x + 44y 12 = 0$

$$a = 1, b = -35, c = -12, f = 22, g = -2,$$
 $h = 1$

Point of intersection is
$$\left(\frac{hf - bg}{ab - h^2}, \frac{gh = af}{ab - h^2}\right)$$

$$= \left(\frac{22-70}{-35-1}, \frac{-2-22}{-35-1}\right) = \left(\frac{-48}{-36}, \frac{-24}{-36}\right) = \left(\frac{4}{3}, \frac{2}{3}\right)$$

Point of intersection of the given lines is $P\left(\frac{4}{3}, \frac{2}{3}\right)$. Given line is 5x + 2y - 8 = 0.

Substituting P in above line,

$$5x + 2y - 8 = 5.\frac{4}{3} + 2.\frac{2}{3} - 8 = \frac{20 + 4 - 24}{3} = 0$$

P lies on the third line 5x + 2y - 8 = 0

- :. The given lines are concurrent.
- 5. Find the distances between the following pairs of parallels straight lines :

i).
$$9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$$

Sol. Given equation is

$$9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$$
.

From above equation a = 9, b=1, c=8, h=-3, g=9, f=-3.

Distance between parallel lines = $2\sqrt{\frac{g^2 - ac}{a(a+b)}}$

$$=2\sqrt{\frac{9^2-9.8}{9(9+1)}}=2\sqrt{\frac{9}{9.10}}=\sqrt{\frac{4}{10}}=\sqrt{\frac{2}{5}}$$

ii.
$$x^2 + 2\sqrt{3}xy + 3y^2 - 3x - 3\sqrt{3}y - 4 = 0$$

ans. $\frac{5}{2}$

- 6. Show that the pairs of lines $3x^2 + 8xy 3y^2 = 0$ and $3x^2 + 8xy 3y^2 + 2x 4y 1 = 0$ form a squares.
- **Sol.** Equation of the first pair of lines is $3x^2 + 8xy 3y^2 = 0$

$$\Rightarrow (x+3y)(3x-y)=0 \qquad \Rightarrow 3x-y=0, x+3y=0$$

Equations of the lines are 3x - y = 0(1) and x + 3y = 0(2)

Equation of the second pair of lines is $3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = 0$

Since
$$3x^2 + 8xy - 3y^2 = (x + 3y)(3x - y)$$

Let
$$3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = (3x - y + c_1)(x + 3y + c_2)$$

Equating the co-efficient of x, we get $c_1 + 3c_2 = 2$

Equation the co-efficient of y, we get $3c_1+c_2=-4$

$$\frac{c_1}{12-2} = \frac{c_2}{-6-4} = \frac{1}{-1-9}$$

$$c_1 = \frac{10}{-10} = -1, c_2 = \frac{-10}{-10} = 1$$

Equations of the lines represented by $3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = 0$ are

$$3x - y - 1 = 0$$
(3)and $x + 3y + 1 = 0$(4)

From above equations, lines (1) and (3) are parallel and lines (2) and (4) are parallel.

Therefore given lines form a parallelogram.

But the adjacent sides are perpendicular, it is a rectangle.(since,(1),(2) are perpendicular and (3),(4) and perpendicular.)

The point of intersection of the pair of lines $3x^2 + 8xy - 3y^2 = 0$ is O(0,0).

Length of the perpendicular from O to (3) =
$$\frac{|0+0+1|}{\sqrt{1+9}} = \frac{1}{\sqrt{10}}$$

Length of the perpendicular from O to (4) =
$$\frac{|0+0+1|}{\sqrt{1+9}} = \frac{1}{\sqrt{10}}$$

Therefore, O is equidistant from lines (3),(4).

Therefore, the distance between the parallel lines is same. Hence the rectangle is a square.

III

1. Find the product of the length of the perpendiculars drawn from (2,1) upon the lines $12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = 0$

Sol. Given pair of lines is
$$12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = 0$$

Now

$$12x^{2} + 25xy + 12y^{2} = 12x^{2} + 16xy + 9xy + 12y^{2}$$
$$= 4x(3x + 4y) + 3y(3x + 4y) = (3x + 4y)(4x + 3y)$$


Let
$$12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = (3x + 4y + c_1)(4x + 3y + c_2)$$

Equating the co-efficient of x, y we get

$$4c_1 + 3c_2 = 10 \Rightarrow 4c_1 + 3c_2 - 10 = 0 \dots (1)$$

$$3c_1 + 4c_2 = 11 \Rightarrow 3c_1 + 4c_2 - 11 = 0 \dots (2)$$

Solving,



$$\frac{c_1}{-33+40} = \frac{c_2}{-30+44} = \frac{1}{16-9}$$

$$c_1 = \frac{7}{7} = 1, c_2 = \frac{14}{7} = 2$$


Therefore given lines are 3x + 4y + 1 = 0 ----(3) and 4x + 3y + 2 = 0 ----(4)

Length of the perpendicular form P(2,1) on $(1) = \frac{6+4+1}{\sqrt{9+16}} = \frac{11}{5}$

Length of the perpendicular from P(2,1) on $(2) = \frac{|8+3+2|}{\sqrt{16+9}} = \frac{13}{5}$

Product of the length of the perpendicular $=\frac{11}{5} \times \frac{13}{5} = \frac{143}{25}$

2. Show that the straight lines $y^2 - 4y + 3 = 0$ and $x^2 + 4xy + 4y^2 + 5x + 10y + 4 = 0$ from a parallelogram and find the lengths of its sides.



Sol. Equation of the first pair of lines is

$$y^2 - 4y + 3 = 0, \Rightarrow (y-1)(y-3) = 0$$

$$\Rightarrow$$
 y-1=0 or y-3=0

 \Rightarrow Equations of the lines are y - 1 = 0(1)

and
$$y - 3 = 0$$
(2)

Equations of (1) and (2) are parallel.

Equation of the second pair of lines is $x^2 + 4xy + 4y^2 + 5x + 10y + 4 = 0$

$$\Rightarrow (x+2y)^2 + 5(x+2y) + 4 = 0$$

$$\Rightarrow (x+2y)^2 + 4(x+2y) + (x+2y) + 4 = 0$$


$$\Rightarrow$$
 $(x+2y)(x+2y+4)+1(x+2y+4)=0$

$$\Rightarrow$$
 $(x+2y+1)(x+2y+4)=0$

$$\Rightarrow$$
 x + 2y + 1 = 0, x + 2y + 4 = 0

Equations of the lines are x+2y+1=0.....(3) and x+2y+4=0.....(4)

Equations of (3) and (4) are parallel .



Solving (1), (3)
$$x + 2 + 1 = 0$$
, $x = -3$

Co-ordinates of A are (-3, 1)

Solving (2), (3)
$$x + 6 + 1 = 0$$
, $x = -7$

Co-ordinates of D are (-7,3)

Solving (1), (4)
$$x + 2 + 4 = 0$$
, $x = -6$

Co-ordinates of B are (-6, 1)

$$AB = \sqrt{(-3+6)^2 + (1-1)^2} = \sqrt{9+0} = 3$$

AD =
$$\sqrt{(-3+7)^2 + (1-3)^2} = \sqrt{16+4} = \sqrt{20} = 2\sqrt{5}$$

Length of the sides of the parallelogram are $3,2\sqrt{5}$

3. Show that the product of the perpendicular distances from the origin to the pair of straight lines represented by $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ is

$$\frac{\left|c\right|}{\sqrt{\left(a-b\right)^{2}+4h^{2}}}$$

Sol. Let
$$l_1 x + m_1 y + n_1 = 0$$
(1)

 $l_2x + m_2y + n_2 = 0$ (2) be the lines represented by

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$$

$$\Rightarrow ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = (l_{1}x + m_{1}y + n_{1})(l_{2}x + m_{2}y + n_{2})$$

$$\Rightarrow l_1 l_2 = a, m_1 m_2 = b, l_1 m_2 + l_2 m_1 = 2h$$

$$l_1 \mathbf{n}_2 + l_2 \mathbf{n}_1 = 2\mathbf{g}, \mathbf{m}_1 \mathbf{n}_2 + \mathbf{m}_2 \mathbf{n}_1 = 2\mathbf{f}, \mathbf{n}_1 \mathbf{n}_2 = \mathbf{c}$$

Perpendicular from origin to (1) = $\frac{|\mathbf{n}_1|}{\sqrt{l_1^2 + \mathbf{m}_1^2}}$

Perpendicular from origin to (2) =
$$\frac{|\mathbf{n}_2|}{\sqrt{l_2^2 + \mathbf{m}_2^2}}$$

Product of perpendiculars

$$= \frac{|\mathbf{n}_{1}|}{\sqrt{l_{1}^{2} + \mathbf{m}_{1}^{2}}} \cdot \frac{|\mathbf{n}_{2}|}{\sqrt{l_{2}^{2} + \mathbf{m}_{2}^{2}}}$$

$$= \frac{|\mathbf{n}_{1}\mathbf{n}_{2}|}{\sqrt{l_{1}^{2}l_{2}^{2} + \mathbf{m}_{1}^{2}\mathbf{m}_{2}^{2} + l_{1}^{2}\mathbf{m}_{2}^{2} + l_{2}^{2}\mathbf{m}_{1}^{2}}}$$

$$= \frac{|\mathbf{n}_{1}\mathbf{n}_{2}|}{\sqrt{(l_{1}l_{2} - \mathbf{m}_{1}\mathbf{m}_{2})^{2} + (l_{1}\mathbf{m}_{2} + l_{2}\mathbf{m}_{1})^{2}}}$$

$$= \frac{|\mathbf{c}|}{\sqrt{(\mathbf{a} - \mathbf{b})^{2} + (2\mathbf{h})^{2}}} = \frac{|\mathbf{c}|}{\sqrt{(\mathbf{a} - \mathbf{b})^{2} + 4\mathbf{h}^{2}}}$$

4. If the equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of intersecting lines, then show that the square of the distance of their point of intersection from the origin is $\frac{c(a+b)-f^2-g^2}{ab-h^2}$. Also show that the square of this distance from origin is $\frac{f^2+g^2}{h^2+h^2}$ if the given lines are perpendicular.

Sol. Let
$$l_1 x + m_1 y + n_1 = 0$$
(1)

$$l_2 x + m_2 y + n_2 = 0$$
(2)

be the lines represented by $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

$$\Rightarrow$$
 ax² + 2hxy + by² + 2gx + 2fy + c

$$= (l_1 x + m_1 y + n_1)(l_2 x + m_2 y + n_2)$$

$$l_1 l_2 = a, m_1 m_2 = b, l_1 m_2 + l_2 m_1 = 2h$$

$$l_1 n_2 + l_2 n_1 = 2g, m_1 n_2 + m_2 n_1 = 2f, n_1 n_2 = c$$
 Solving (1) and (2)

$$\frac{\mathbf{x}}{\mathbf{m}_1 \mathbf{n}_2 - \mathbf{m}_2 \mathbf{n}_1} = \frac{\mathbf{y}}{l_2 \mathbf{n}_1 - l_1 \mathbf{n}_2} = \frac{1}{l_1 \mathbf{m}_2 - l_2 \mathbf{m}_2}$$

The point of intersection is P=
$$\left[\frac{m_1 n_2 - m_2 n_1}{l_1 m_2 - l_2 m_1}, \frac{l_2 n_1 - l_1 n_2}{l_1 m_2 - l_2 m_1}\right]$$

$$OP^{2} = \frac{\left(m_{1}n_{2} - m_{2}n_{1}\right)^{2} + \left(l_{2}n_{1} - l_{1}n_{2}\right)^{2}}{\left(l_{1}m_{2} - l_{2}m_{1}\right)^{2}}$$

$$=\frac{\left(m_{1}n_{2}+m_{2}n_{1}\right)^{2}-4m_{1}m_{2}n_{1}n_{2}+\left(l_{1}n_{2}+l_{2}n_{1}\right)^{2}-4l_{1}l_{2}n_{1}n_{2}}{\left(l_{1}m_{2}+l_{2}m_{1}\right)^{2}-4l_{1}l_{2}m_{1}m_{2}}$$

$$=\frac{4f^2 - 4abc + 4g^2 - 4ac}{4h^2 - 4ab}$$

$$=\frac{c(a+b)-f^2-g^2}{ab-h^2}.$$

If the given pair of lines are perpendicular, then $a + b = 0 \Rightarrow a = -b$


$$\Rightarrow OP^{2} = \frac{0 - f^{2} - g^{2}}{(-b)b - h^{2}} = \frac{f^{2} + g^{2}}{h^{2} + b^{2}}$$

HOMOGENISATION

THEOREM

The equation to the pair of lines joining the origin to the points of intersection of the curve $S = ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ and the line

$$L = lx + my + n = 0 \text{ is } ax^2 + 2hxy + by^2 + (2gx + 2fy) \left(\frac{lx + my}{-n}\right) + c\left(\frac{lx + my}{-n}\right)^2 = 0 - - (1)$$



Eq (1) represents the combined equation of the pair of lines \overline{OA} and \overline{OB} .

I

- 1. Find the equation of the lines joining the origin to the point of intersection of $x^2 + y^2 = 1$ and x + y = 1
- **Sol.** The given curves are $x^2 + y^2 = 1$(1)

$$x + y = 1$$
(2)

Homogenising (1) with the help of (2) then $x^2 + y^2 = 1^2$

$$\Rightarrow x^2 + y^2 = (x + y)^2 = x^2 + y^2 + 2xy$$
 i.e. $2xy = 0 \Rightarrow xy = 0$

- 2. Find the angle between the lines joining the origin to points of intersection of $y^2 = x$ and x + y = 1.
- **Sol.** Equation of the curve is $y^2 = x$ (1) and Equation of line is x + y = 1(2)

Harmogonsing (1) with the help of (2)


$$Y^2 - x \cdot 1 = 0 \implies y^2 = x(x + y) = x^2 + xy$$

 \Rightarrow $x^2 + xy - y^2 = 0$ which represents a pair of lines. From this equation a + b = 1 - 1 = 0

The angle between the lines is 90° .

II

1. Show that the lines joining the origin to the points of intersection of the curve $x^2 - xy + y^2 + 3x + 3y - 2 = 0$ and the straight line $x - y - \sqrt{2} = 0$ are mutually perpendicular.



Sol.

Let A,B the the points of intersection of the line and the curve.

Equation of the curve is $x^2 - xy + y^2 + 3x + 3y - 2 = 0$ (1)

Equation of the line AB is $x - y - \sqrt{2} = 0$

$$\Rightarrow x - y = \sqrt{2} \Rightarrow \frac{x - y}{\sqrt{2}} = 1 \qquad \dots (2)$$

Homogenising, (1) with the help of (2) combined equation of OA, OB is

$$x^2 - xy + y^2 + 3x \cdot 1 + 3y \cdot 1 - 2 \cdot 1^2 = 0$$

$$\Rightarrow x^{2} - xy + y^{2} + 3(x + y)\frac{x - y}{\sqrt{2}} - 2\frac{(x - y)^{2}}{2} = 0$$

$$\Rightarrow x^{2} - xy + y^{2} + \frac{3}{\sqrt{2}}(x^{2} - y^{2}) - (x^{2} - 2xy + y^{2}) = 0$$

$$\Rightarrow x^{2} - xy + y^{2} + \frac{3}{\sqrt{2}}x^{2} - \frac{3}{\sqrt{2}}y^{2} - x^{2} + 2xy - y^{2} = 0$$


$$\Rightarrow \frac{3}{\sqrt{2}}x^2 + xy - \frac{3}{\sqrt{2}}y^2 = 0$$

$$\Rightarrow a+b=\frac{3}{\sqrt{2}}-\frac{3}{\sqrt{2}}=0$$

: OA, OB are perpendicular.

- 2. Find the values of k, if the lines joining the origin to the points of intersection of the curve $2x^2 2xy + 3y^2 + 2x y 1 = 0$ and the line x + 2y = k are mutually perpendicular.
- **Sol.** Given equation of the curve is $S = 2x^2 2xy + 3y^2 + 2x y 1 = 0 \dots (1)$

Equation of AB is x + 2y = k



$$\frac{x+2y}{k} = 1 \qquad \dots (2$$

Le t A,B the the points of intersection of the line and the curve.

Homogenising, (1) with the help of (2), the combined equation of OA,OB is

$$2x^2 - 2xy + 3y^2 + 2x \cdot 1 - y \cdot 1 - 1^2 = 0$$

$$2x^{2} - 2xy + 3y^{2} + 2x\frac{(x+2y)}{k} - y\frac{(x+2y)}{k} = \frac{(x+2y)^{2}}{k^{2}} = 0$$

$$\Rightarrow 2k^2x^2 - 2k^2xy + 3k^2y^2 + 2kx(x+2y) - ky(x+2y) - (x+2y)^2 = 0$$

$$\Rightarrow 2k^2x^2 - 2k^2xy + 3k^2y^2 + 2kx^2 + 4kxy - kxy - 2ky^2 - x^2 - 4xy - 4y^2 = 0$$

$$\Rightarrow (2k^2 + 2k - 1)x^2 + (-2k^2 + 3k - 4)xy + (3k^2 - 2k - 4)y^2 = 0$$


Given that above lines are perpendicular, Co-efficient x^2 + co-efficient of y^2 =0

$$\Rightarrow 2k^2 + 2k - 1 + 3k^2 - 2k - 4 = 0$$

$$\Rightarrow 5k^2 = 5 \Rightarrow k^2 = 1 : k = \pm 1$$

3. Find the angle between the lines joining the origin to the points of intersection of the curve $x^2 + 2xy + y^2 + 2x + 2y - 5 = 0$ and the line 3x - y + 1 = 0

Sol.



Equation of the curve is

$$x^{2} + 2xy + y^{2} + 2x + 2y - 5 = 0$$
....(1)

Equation of AB is
$$3x - y + 1 = 0 \Rightarrow y - 3x = 1$$
(2

Let A,B the the points of intersection of the line and the curve.

Homogenising (1) with the help of (2), combined equation of OA, OB is

$$x^{2} + 2xy + y^{2} + 2x.1 + 2y.1 - 5.1^{2} = 0$$

$$\Rightarrow x^{2} + 2xy + y^{2} + 2x(y - 3x) + 2y(y - 3x) - 5(y - 3x)^{2} = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + 2xy - 6x^2 + 2y^2 - 6xy - 5(y^2 + 9x^2 - 6xy) = 0$$

$$\Rightarrow -5x^2 - 2xy + 3y^2 - 5y^2 - 45x^2 + 30xy = 0$$

$$\Rightarrow -50x^2 + 28xy - 2y^2 = 0 \Rightarrow 25x^2 - 14xy + y^2 = 0$$


let θ be the angle between OA and OB, then

$$\cos \theta = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|25+1|}{\sqrt{(25-1)^2 + 196}} = \frac{26}{\sqrt{576 + 196}} = \frac{26}{\sqrt{772}}$$

$$= \frac{26}{2\sqrt{193}} = \frac{13}{\sqrt{193}} :: \theta = \cos^{-1}\left(\frac{13}{\sqrt{193}}\right)$$

1. Find the condition for the chord lx + my = 1 of the circle $x^2 + y^2 = a^2$ (whose centre is the origin) to subtend a right angle at the origin.

Sol.



Equation of the circle $x^2 + y^2 = a^2$ (1)

Equation of AB is lx + my = 1(2)

Let A,B the the points of intersection of the line and the curve

Homogenising (1) with the help of (2) ,the combined equation of OA, OB is

$$x^{2} + y^{2} = a^{2}.1^{2} \Rightarrow x^{2} + y^{2} = a^{2} (lx + my)^{2}$$

$$= a^{2} (l^{2}x^{2} + m^{2}y^{2} + 2lmxy) = a^{2}l^{2}x^{2} + a^{2}m^{2}y^{2} + 2a^{2}lmxy$$

$$\Rightarrow a^2 l^2 x^2 + 2a^2 l m x y + a^2 m^2 y^2 - x^2 - y^2 = 0$$


$$\Rightarrow (a^{2}l^{2}-1)x^{2}+2a^{2}lmxy+(a^{2}m^{2}-1)y^{2}=0$$

Since OA, OB are perpendicular, Coefficient of x^2 + co-efficient of y^2 =0

$$\Rightarrow$$
 $a^2l^2 - 1 + a^2m^2 - 1 = 0 \Rightarrow a^2(l^2 + m^2) = 2$ which is the required condition

2. Find the condition for the lines joining the origin to the points of intersection of the circle $x^2 + y^2 = a^2$ and the line lx + my = 1 to coincide.

Sol.



Equation of the circle is $x^2 + y^2 = a^2$ (1)

Equation of AB is $lx + my = 1 \dots (2)$.

Le t A,B the the points of intersection of the line and the curve.

Homogenising (1) with the help of (2),

Then the combined equation of OA, OB is $x^2 + y^2 = a^2 \cdot 1^2$

$$x^{2} + y^{2} = a^{2} (lx + my)^{2} = a^{2} (l^{2}x^{2} + m^{2}y^{2} + 2lmxy)$$

$$\Rightarrow x^2 + y^2 = a^2 l^2 x^2 + a^2 m^2 y^2 + 2a^2 l m x y$$

$$\Rightarrow (a^{2}l^{2}-1)x^{2}+2a^{2}lmxy+(a^{2}m^{2}-1)y^{2}=0$$

Since OA, OB are coincide $\Rightarrow h^2 = ab$

$$\Rightarrow a^4 l^2 m^2 = (a^2 l^2 - 1)(a^2 m^2 - 1) \implies a^4 l^2 m^2 = a^4 l^2 m^2 - a^2 l^2 - a^2 m^2 + 1$$

$$\therefore a^2 l^2 - a^2 m^2 + 1 = 0 \implies a^2 (l^2 + m^2) = 1$$

This is the required condition.

- 3. Write down the equation of the pair of straight lines joining the origin to the points of intersection of the lines 6x y + 8 = 0 with the pair of straight lines $3x^2 + 4xy 4y^2 11x + 2y + 6 = 0$. Show that the lines so obtained make equal angles with the coordinate axes.
- **Sol.** Given pair of line is $3x^2 + 4xy 4y^2 11x + 2y + 6 = 0$...(1)

Given line is
$$6x - y + 8 = 0 \Rightarrow \frac{6x - y}{-8} = 1 \Rightarrow \frac{y - 6x}{8} = 1 - - - (2)$$

Homogenising (1) w.r.t (2)

$$3x^{2} + 4xy - 4y^{2} - (11x - 2y)\left(\frac{y - 6x}{8}\right) + 6\left(\frac{y - 6x}{8}\right)^{2} = 0$$

$$64 \left\lceil 3x^2 + 4xy - 4y^2 \right\rceil - 8 \left\lceil 11xy - 66x^2 - 2y^2 + 12xy \right\rceil + 6 \left[y^2 + 36x^2 - 12xy \right] = 0$$

$$\Rightarrow$$
 936x² + 256xy - 256xy - 234y² = 0

$$\Rightarrow 468x^2 - 117y^2 = 0$$

$$\Rightarrow 4x^2 - y^2 = 0 - - (3)$$

Is eq. of pair of lines joining the origin to the point of intersection of (1) and (2).

The eq. pair of angle bisectors of (3) is

$$h(x^2 - y^2) - (a - b)xy = 0$$

$$\Rightarrow 0(x^2-y^2)-(4-1)xy=0 \Rightarrow xy=0$$

x = 0 or y = 0 which are the eqs. is of co-ordinates axes

:. The pair of lines are equally inclined to the co-ordinate axes

- 4. If the straight lines given by $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ intersect on Y-axis, show that $2fgh bg^2 ch^2 = 0$
- **Sol.** Given pair of lines is $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

Equation of Y-axis is x = 0 then equation becomes $by^2 + 2fy + c = 0$ (1)

Since the given pair of lines intersect on Y – axis, the roots or equation (1) are equal.

 \therefore Discriminate = 0

$$\Rightarrow (2f)^2 - 4.b.c = 0 \Rightarrow 4f^2 - 4bc = 0$$

$$\Rightarrow$$
 f² - bc = 0 \Rightarrow f² = bc

Since the given equation represents a pair of lines


$$abc + 2fgh + af^{2} - bg^{2} - ch^{2} = 0$$

$$\Rightarrow a(f^2) + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow$$
 2fgh - bg² - ch² = 0

- 5. Prove that the lines represented by the equations $x^2 4xy + y^2 = 0$ and x + y = 3 form an equilateral triangle.
- **Sol.** Since the straight line L: x + y = 3 makes 45° with the negative direction of the X –axis, none of the lines which makes 60° with the line L is vertical. If 'm' is the slope of one such straight line, then $\sqrt{3} = \tan 60^{\circ} = \left| \frac{m+1}{1-m} \right|$ and so, satisfies the equation $(m+1)^2 = 3(m-1)^2$

Or
$$m^2 - 4m + 1 = 0$$
(1)



But the straight line having slope 'm' and passing through the origin is

$$y = mx$$
(2)

So the equation of the pair of lines passing through the origin and inclined at 60° with the line L is obtained by eliminating 'm' from the equations (1) and (2).

Therefore the combined equation of this pair of lines is $\left(\frac{y}{x}\right)^2 - 4\left(\frac{y}{x}\right) + 1 = 0$ (i.e,)

$$x^2 - 4xy + y^2 = 0$$

Which is the same as the given pair of lines. Hence, the given traid of lines form an equilateral triangle.

6. Show that the product of the perpendicular distances from a point (α, β) to

the pair of straight lines
$$ax^2 + 2hxy + by^2 = 0$$
 is $\frac{\left|a\alpha^2 + 2h\alpha\beta + b\beta^2\right|}{\sqrt{\left(a-b\right)^2 + 4h^2}}$

Sol. Let
$$ax^2 + 2hxy + by^2 \equiv (l_1x + m_1y)(l_2x + m_2y)$$

Then the separate equations of the lines represented by the equation

$$ax^2 + 2hxy + by^2 = 0$$
 are $L_1: l_1x + m_1y = 0$ and $L_2: l_2x + m_2y = 0$

Also, we have
$$l_1 l_2 = a$$
; $m_1 m_2 = b$ and $l_1 m_2 + l_2 m_1 = 2h$

$$d_1$$
 =length of the perpendicular from (α, β) to $L_1 = \frac{|l_1\alpha + m_1\beta|}{\sqrt{l_1^2 + m_1^2}}$

$$d_2$$
 =length of the perpendicular from (α, β) to L_2 $L_2 = \frac{|l_2\alpha + m_2\beta|}{\sqrt{l_2^2 + m_2^2}}$

Then, the product of the lengths of the perpendiculars from (α, β) to the given pair of lines = d_1d_2

$$= \frac{\left| (l_1 \alpha + m_1 \beta) (l_2 \alpha + m_2 \beta) \right|}{\sqrt{(l_1^2 + m_1^2)(l_2^2 + m_2^2)}} = \frac{\left| a\alpha^2 + 2h\alpha\beta + b\beta^2 \right|}{\sqrt{(a-b)^2 + 4h^2}}$$

PROBLEMS FOR PRACTICE.

- 1. If the lines xy+x+y+1=0 and x+ay-3=0 are concurrent, find a.
- 2. The equation $ax^2 + 3xy 2y^2 5x + 5y + c = 0$ represents two straight lines perpendicular to each other. Find a and c.
- 3. Find λ so that $x^2 + 5xy + 4y^2 + 3x + 2y + \lambda = 0$ may represent a pair of straight lines. Find also the angle between them for this value of λ .
- 4. If $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents the straight lines equidistant from the origin, show that $f^4 g^4 = c(bf^2 ag^2)$
- 5. Find the centroid of the triangle formed by the lines $12x^2 20xy + 7y^2 = 0$ and 2x - 3y + 4 = 0

ANS. =
$$\left(\frac{8}{3}, \frac{8}{3}\right)$$

- 6. Let $aX^2 + 2hXY + bY^2 = 0$ represent a pair of straight lines. Then show that the equation of the pair of straight lines.
 - i)Passing through (x_o, y_o) and parallel to the given pair of lines is

$$a(x-x_o)^2+2h(x-x_o)(y-y_o)+b(y-y_o)^2=0$$
 ii) Passing through (x_o,y_o) and perpendicular to the given pair of lines

is
$$b(x-x_o)^2 - 2h(x-x_o)(y-y_o) + a(y-y_o)^2 = 0$$

- 7. Find the angle between the straight lines represented by $2x^2 + 3xy 2y^2 5x + 5y 3 = 0$
- 8. Find the equation of the pair of lines passing through the origin and perpendicular to the pair of lines $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$
- 9. If $x^2 + xy + 2y^2 + 4x y + k = 0$ represents a pair of straight lines find k.
- 10. Prove that equation $2x^2 + xy 6y^2 + 7y 2 = 0$ represents a pair of straight line.
- 11. Prove that the equation $2x^2+3xy-2y^2-x+3y-1=0$ represents a pair of perpendicular straight lines.
- 12. Show that the equation $2x^2-13xy-7y^2+x+23y-6=0$ represents a pair of straight lines. Also find the angle between the co-ordinates of the point of intersection of the lines.
- 13. Find that value of λ for which the equation $\lambda x^2 10xy + 12y^2 + 5x 16y 3 = 0$ represents a pair of straight lines.
- 14. Show that the pair of straight lines $6x^2-5xy-6y^2=0$ and $6x^2-5xy-6y^2+x+5y-1=0$ form a square.
- 15. Show that the equation $8x^2 24xy + 18y^2 6x + 9y 5 = 0$ represents pair of parallel straight lines are find the distance between them.