

## PAIR OF LINES-SECOND DEGREE GENERAL EQUATION

### THEOREM

If the equation  $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents a pair of straight lines then

$$\text{i) } \Delta \equiv abc + 2fgh - af^2 - bg^2 - ch^2 = 0 \quad \text{and (ii) } h^2 \geq ab, g^2 \geq ac, f^2 \geq bc$$

### Proof:

Let the equation  $S = 0$  represent the two lines  $l_1x + m_1y + n_1 = 0$  and  $l_2x + m_2y + n_2 = 0$ .

Then

$$\begin{aligned} & ax^2 + 2hxy + by^2 + 2gx + 2fy + c \\ & \equiv (l_1x + m_1y + n_1)(l_2x + m_2y + n_2) = 0 \end{aligned}$$

Equating the co-efficients of like terms, we get

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b, \text{ and } l_1n_2 + l_2n_1 = 2g, m_1n_2 + m_2n_1 = 2f, n_1n_2 = c$$

(i) Consider the product  $(2h)(2g)(2f)$

$$\begin{aligned} & = (l_1m_2 + l_2m_1)(l_1n_2 + l_2n_1)(m_1n_2 + m_2n_1) \\ & = l_1l_2(m_1^2n_2^2 + m_2^2n_1^2) + m_1m_2(l_1^2n_2^2 + l_2^2n_1^2) + n_1n_2(l_1^2m_2^2 + l_2^2m_1^2) + 2l_1l_2m_1m_2n_1n_2 \\ & = l_1l_2[(m_1n_2 + m_2n_1)^2 - 2m_1m_2n_1n_2] + m_1m_2[(l_1n_2 + l_2n_1)^2 - 2l_1l_2n_1n_2] \\ & \quad + n_1n_2[(l_1m_2 + l_2m_1)^2 - 2l_1l_2m_1m_2] + 2l_1l_2m_1m_2n_1n_2 \\ & = a(4f^2 - 2bc) + b(4g^2 - 2ac) + c(4h^2 - 2ab) \\ & \quad 8fgh = 4[af^2 + bg^2 + ch^2 - abc] \\ & \therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0 \end{aligned}$$

$$\begin{aligned} \text{ii) } h^2 - ab & = \left( \frac{l_1m_2 + l_2m_1}{2} \right)^2 - l_1l_2m_1m_2 = \frac{(l_1m_2 + l_2m_1)^2 - 4l_1l_2m_1m_2}{4} \\ & = \frac{(l_1m_2 - l_2m_1)^2}{4} \geq 0 \end{aligned}$$

Similarly we can prove  $g^2 \geq ac$  and  $f^2 \geq bc$

### NOTE :

If  $\Delta = abc + 2fgh - af^2 - bg^2 - ch^2 = 0$ ,  $h^2 \geq ab$ ,  $g^2 \geq ac$  and  $f^2 \geq bc$ , then the equation  $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents a pair of straight lines

## CONDITIONS FOR PARALLEL LINES-DISTANCE BETWEEN THEM

### THEOREM

If  $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents a pair of parallel lines then  $h^2 = ab$  and  $bg^2 = af^2$ . Also the distance between the two parallel lines is

$$2\sqrt{\frac{g^2 - ac}{a(a+b)}} \quad (\text{or}) \quad 2\sqrt{\frac{f^2 - bc}{b(a+b)}}$$

### Proof :

Let the parallel lines represented by  $S = 0$  be  
 $lx + my + n_1 = 0$  -- (1)  $lx + my + n_2 = 0$  -- (2)

$$\therefore ax^2 + 2hxy + 2gx + 2fy + c$$

$$\equiv (lx + my + n_1)(lx + my + n_2)$$

Equating the like terms

$$l^2 = a \quad \text{-- (3)} \quad 2lm = 2h \quad \text{-- (4)}$$

$$m^2 = b \quad \text{-- (5)} \quad l(n_1 + n_2) = 2g \quad \text{-- (6)}$$

$$m(n_1 + n_2) = 2f \quad \text{-- (7)} \quad n_1n_2 = c \quad \text{-- (8)}$$

From (3) and (5),  $l^2m^2 = ab$  and from (4)  $h^2 = ab$ .

$$\text{Dividing (6) and (7)} \quad \frac{l}{m} = \frac{g}{f} \Rightarrow \frac{l^2}{m^2} = \frac{g^2}{f^2},$$

$$\therefore \frac{a}{b} = \frac{g^2}{f^2} \Rightarrow bg^2 = af^2$$

Distance between the parallel lines (1) and (2) is

$$\begin{aligned} &= \left| \frac{n_1 - n_2}{\sqrt{l^2 + m^2}} \right| = \frac{\sqrt{(n_1 + n_2)^2 - 4n_1n_2}}{\sqrt{l^2 + m^2}} \\ &= \frac{\sqrt{(4g^2 / l^2) - 4c}}{\sqrt{a+b}} \quad \text{or} \quad \frac{\sqrt{(4f^2 / m^2) - 4c}}{\sqrt{a+b}} \\ &= 2\sqrt{\frac{g^2 - ac}{a(a+b)}} \quad (\text{or}) \quad 2\sqrt{\frac{f^2 - bc}{b(a+b)}} \end{aligned}$$

## POINT OF INTERSECTION OF PAIR OF LINES THEOREM

The point of intersection of the pair of lines represented by

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0 \text{ when } h^2 > ab \text{ is } \left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$$

**Proof:**

Let the point of intersection of the given pair of lines be  $(x_1, y_1)$ . Transfer the origin to  $(x_1, y_1)$  without changing the direction of the axes.

Let  $(X, Y)$  represent the new coordinates of  $(x, y)$ . Then  $x = X + x_1$  and  $y = Y + y_1$ .

Now the given equation referred to new axes will be

$$\begin{aligned} a(X + x_1)^2 + 2h(X + x_1)(Y + y_1) + b(Y + y_1)^2 + 2g(X + x_1) + 2f(Y + y_1) + c &= 0 \\ \Rightarrow aX^2 + 2hXY + bY^2 + 2X(ax_1 + hy_1 + g) + 2Y(hx_1 + by_1 + f) \\ + (ax_1^2 + 2hx_1y_1 + by_1^2 + 2gx_1 + 2fy_1 + c) &= 0 \end{aligned}$$

Since this equation represents a pair of lines passing through the origin it should be a homogeneous second degree equation in  $X$  and  $Y$ . Hence the first degree terms and the constant term must be zero. Therefore,

$$ax_1 + hy_1 + g = 0 \quad \text{-- (1)}$$

$$hx_1 + by_1 + f = 0 \quad \text{-- (2)}$$

$$ax_1^2 + 2hx_1y_1 + by_1^2 + 2gx_1 + 2fy_1 + c = 0 \quad \text{-- (3)}$$

But (3) can be rearranged as

$$x_1(ax_1 + hy_1 + g) + y_1(hx_1 + by_1 + f) + (gx_1 + fy_1 + c) = 0$$

$$\Rightarrow gx_1 + fy_1 + c = 0 \quad \text{--(4)}$$

Solving (1) and (2) for  $x_1$  and  $y_1$

$$\frac{x_1}{hf - bg} = \frac{y_1}{gh - af} = \frac{1}{ab - h^2}$$

$$\therefore x_1 = \frac{hf - bg}{ab - h^2} \text{ and } y_1 = \frac{gh - af}{ab - h^2}$$

Hence the point of intersection of the given pair of lines is  $\left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

## THEOREM


If the pair of lines  $ax^2 + 2hxy + by^2 = 0$  and the pair of lines

$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  form a rhombus then  $(a-b)fg + h(f^2 - g^2) = 0$ .

### Proof:

The pair of lines  $ax^2 + 2hxy + by^2 = 0$  -- (1) is parallel to the lines

$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  -- (2)



Now the equation

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c + \lambda(ax^2 + 2hxy + by^2) = 0$$

Represents a curve passing through the points of intersection of (1) and (2).

Substituting  $\lambda = -1$ , in (3) we obtain  $2gx + 2fy + c = 0$  ... (4) Equation (4) is a straight line passing through A and B and it is the diagonal  $\overline{AB}$

The point of intersection of (2) is  $C = \left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

$$\Rightarrow \text{Slope of } \overline{OC} = \frac{gh - af}{hf - bg}$$

In a rhombus the diagonals are perpendicular  $\Rightarrow (\text{Slope of } \overline{OC})(\text{Slope of } \overline{AB}) = -1$

$$\Rightarrow \left( \frac{gh - af}{hf - bg} \right) \left( -\frac{g}{f} \right) = -1$$

$$\Rightarrow g^2h - afg = hf^2 - bfg$$

$$\Rightarrow (a-b)fg + h(f^2 - g^2) = 0$$


$$\frac{g^2 - f^2}{a-b} = \frac{fg}{h}$$

## THEOREM

If  $ax^2 + 2hxy + by^2 = 0$  be two sides of a parallelogram and  $px + qy = 1$  is one diagonal, then the other diagonal is  $y(bp - hq) = x(aq - hp)$

**proof:**

Let  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  be the points where the diagonal



$px + qy = 1$  meets the pair of lines.

$\overline{OR}$  and  $\overline{PQ}$  bisect each other at  $M(\alpha, \beta)$ .

$$\therefore \alpha = \frac{x_1 + x_2}{2} \text{ and } \beta = \frac{y_1 + y_2}{2}$$

Eliminating  $y$  from  $ax^2 + 2hxy + by^2 = 0$  -- (1)

and  $px + qy = 1$  -- (2)

$$ax^2 + 2hx\left(\frac{1-px}{q}\right) + b\left(\frac{1-px}{q}\right)^2 = 0$$

$$\Rightarrow x^2(aq^2 - 2hpq + bp^2) + 2x(hp - bp) + b = 0$$

The roots of this quadratic equation are  $x_1$  and  $x_2$  where

$$x_1 + x_2 = -\frac{2(hq - bp)}{aq^2 - 2hpq - bp^2}$$

$$\Rightarrow \alpha = \frac{(bp - hq)}{(aq^2 - 2hpq + bp^2)}$$

Similarly by eliminating  $x$  from (1) and (2) a quadratic equation in  $y$  is obtained and  $y_1,$

$y_2$  are its roots where

$$y_1 + y_2 = -\frac{2(hp - aq)}{aq^2 - 2hpq - bp^2} \Rightarrow \beta = \frac{(aq - hp)}{(aq^2 - 2hpq + bp^2)}$$

Now the equation to the join of  $O(0, 0)$  and  $M(\alpha, \beta)$  is  $(y-0)(0-\alpha) = (x-0)(0-\beta)$

$$\Rightarrow \alpha y = \beta x$$

Substituting the values of  $\alpha$  and  $\beta$ , the equation of the diagonal OR

$$\text{is } y(bp - hq) = x(aq - hp) .$$

### EXERCISE 4B

#### I

1. Find the angle between the lines represented by  $2x^2 + xy - 6y^2 + 7y - 2 = 0$ .

**Sol.** Given equation is

$$2x^2 + xy - 6y^2 + 7y - 2 = 0 \text{ Comparing with}$$

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0 \text{ then}$$

$$a = 2, b = -6, c = -2, g = 0, f = \frac{7}{2}, h = \frac{1}{2}$$

Angle between the lines is given by

$$\cos \alpha = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|2-6|}{\sqrt{(2+6)^2 + 1}} = \frac{4}{\sqrt{65}} \Rightarrow \alpha = \cos^{-1} \left( \frac{4}{\sqrt{65}} \right)$$

2. Prove that the equation  $2x^2 + 3xy - 2y^2 + 3x + y + 1 = 0$  represents a pair of perpendicular lines.

**Sol.** From given equation  $a = 2, b = -2$  and  $a + b = 2 + (-2) = 0$

$$\Rightarrow \text{angle between the lines is } 90^\circ. \quad \therefore \text{The given lines are perpendicular.}$$

## II

1. Prove that the equation  $3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$  represents a pair of straight lines and find the co-ordinates of the point of intersection.

**Sol.** The given equation is  $3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$

Comparing with  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ , we get

$$a = 3, \quad b = 2, \quad c = 2, \quad 2f = 5 \quad \Rightarrow f = \frac{5}{2}$$

$$2g = 5 \Rightarrow g = \frac{5}{2}, \quad 2h = 7 \Rightarrow h = \frac{7}{2}$$

$$\Delta = abc + 2fgh - af^2 - bg^2 - ch^2$$

$$= 3(2)(2) + 2 \cdot \frac{5}{2} \cdot \frac{5}{2} \cdot \frac{7}{2} - 3 \cdot \frac{25}{4} - 2 \cdot \frac{25}{4} - 2 \cdot \frac{49}{4}$$

$$= \frac{1}{4}(48 + 175 - 75 - 50 - 98)$$

$$= \frac{1}{2}(223 - 223) = 0$$

$$h^2 - ab = \left(\frac{7}{2}\right)^2 - 3 \cdot 2 = \frac{49}{4} - 6 = \frac{25}{4} > 0$$

$$f^2 - bc = \left(\frac{5}{2}\right)^2 - 2 \cdot 2 = \frac{25}{4} - 4 = \frac{9}{4} > 0$$

$$g^2 - ac = \left(\frac{5}{2}\right)^2 - 3 \cdot 2 = \frac{25}{4} - 6 = \frac{1}{4} > 0$$

$\therefore$  The given equation represents a pair of lines.

The point of intersection of the lines is  $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$

$$= \left( \frac{\frac{7}{2} \cdot \frac{5}{2} - 2 \cdot \frac{5}{2}}{6 - \frac{49}{4}}, \frac{\frac{5}{2} \cdot \frac{7}{2} - 3 \cdot \frac{5}{2}}{6 - \frac{49}{4}} \right) = \left( \frac{35 - 20}{24 - 29}, \frac{35 - 30}{24 - 49} \right)$$

$$= \left( \frac{+15}{-25}, \frac{5}{-28} \right) = \left( \frac{-3}{5}, -\frac{1}{5} \right)$$

Point of intersection is  $p \left( \frac{-3}{5}, -\frac{1}{5} \right)$

- 2. Find the value of k, if the equation  $2x^2 + kxy - 6y^2 + 3x + y + 1 = 0$  represents a pair of straight lines. Find the point of intersection of the lines and the angle between the straight lines for this value of k.**

**Sol.** The given equation is  $2x^2 + kxy - 6y^2 + 3x + y + 1 = 0$

$$a = 2, b = -6, c = 1, f = \frac{1}{2}, 2g = 3 \Rightarrow g = \frac{3}{2}, h = \frac{k}{2}$$

Since the given equation is representing a pair of straight lines, therefore

$$\Delta = abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow -12 + 2 \cdot \frac{1}{2} \cdot \frac{3}{2} \cdot \left( +\frac{k}{2} \right) - 2 \cdot \frac{1}{4} + 6 \cdot \frac{9}{4} - \frac{k^2}{4} = 0$$

$$\Rightarrow -48 + 3k - 2 + 54 - k^2 = 0$$

$$\Rightarrow -k^2 + 3k + 4 = 0 \Rightarrow k^2 - 3k - 4 = 0$$

$$\Rightarrow (k - 4)(k + 1) = 0$$

$$\Rightarrow k = 4 \text{ or } -1$$

**Case (i)  $k = -1$**

Point of intersection is  $\left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$



$$\left( \frac{+\frac{1}{2} \cdot \frac{1}{2} + 6 \cdot \frac{3}{2}}{-12 - \frac{1}{4}}, \frac{\frac{3}{2} \left( -\frac{1}{2} \right) - 2 \cdot \frac{1}{2}}{-12 - \frac{1}{4}} \right) = \left( \frac{-1+36}{-49}, \frac{-3-4}{-49} \right)$$

$$= \left( \frac{35}{-49}, \frac{-7}{-49} \right) = \left( \frac{-5}{7}, \frac{1}{7} \right)$$

Point of intersection is  $\left( \frac{-5}{7}, \frac{1}{7} \right)$

$$\text{Angle between the lines} = \cos \alpha = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|2-6|}{\sqrt{(2+6)^2 + 4}} = \left( \frac{4}{\sqrt{65}} \right)$$

**Case (ii)**  $k = 4$

$$\left( \frac{2 \cdot \frac{1}{2} + 6 \cdot \frac{3}{2}}{-12 - 4}, \frac{\frac{3}{2} \cdot 2 - 2 \cdot \frac{1}{2}}{-12 - 4} \right) = \left( -\frac{5}{8}, -\frac{1}{8} \right)$$

Point of intersection is  $P \left( -\frac{5}{8}, -\frac{1}{8} \right)$  and angle between the lines is

$$\begin{aligned} \cos \alpha &= \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} \\ &= \frac{|2-6|}{\sqrt{(2+6)^2 + 16}} = \frac{4}{4\sqrt{5}} = \frac{1}{\sqrt{5}} \end{aligned}$$

$$\alpha = \cos^{-1} \left( \frac{1}{\sqrt{5}} \right)$$

3. Show that the equation  $x^2 - y^2 - x + 3y - 2 = 0$  represents a pair of perpendicular lines and find their equations.

Sol. Given equation is  $x^2 - y^2 - x + 3y - 2 = 0 \Rightarrow a = 1, b = 1, c = -2, f = \frac{3}{2}, g = -\frac{1}{2},$

$$h = 0$$

$$\text{Now } \Delta = abc + 2fgh - af^2 - bg^2 - ch^2$$

$$= 1(-1)(-2) + 0 - 1 \cdot \frac{9}{4} + 1 \cdot \frac{1}{4} + 0 = +2 - \frac{9}{4} + \frac{1}{4} = 0$$

$$h^2 - ab = 0 - 1(-1) = 1 > 0$$

$$f^2 - bc = \frac{9}{4} - 2 = \frac{1}{4} > 0$$

$$g^2 - ac = \frac{1}{4} + 2 = \frac{9}{4} > 0$$

$$\text{And } a + b = 1 - 1 = 0$$

The given equation represent a pair of perpendicular lines.

$$\text{Let } x^2 - y^2 - x + 3y - 2 = (x + y + c_1)(x - y + c_2)$$

$$\text{Equating the coefficients of } x \Rightarrow c_1 + c_2 = -1$$

$$\text{Equating the co-efficient of } y \Rightarrow -c_1 + c_2 = 3$$

$$\text{Adding } 2c_2 = 2 \Rightarrow c_2 = 1$$

$$c_1 + c_2 = -1 \Rightarrow c_1 + 1 = -1, c_1 = -2$$

Equations of the lines are  $x + y - 2 = 0$  and  $x - y + 1 = 0$

4. Show that the lines  $x^2 + 2xy - 35y^2 - 4x + 44y - 12 = 0$  and  $5x + 2y - 8 = 0$  are concurrent.

**Sol.** Equation of the given lines are  $x^2 + 2xy - 35y^2 - 4x + 44y - 12 = 0$

$$a = 1, b = -35, c = -12, f = 22, g = -2, \quad h = 1$$

Point of intersection is  $\left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

$$= \left( \frac{22 - 70}{-35 - 1}, \frac{-2 - 22}{-35 - 1} \right) = \left( \frac{-48}{-36}, \frac{-24}{-36} \right) = \left( \frac{4}{3}, \frac{2}{3} \right)$$

Point of intersection of the given lines is  $P\left(\frac{4}{3}, \frac{2}{3}\right)$ . Given line is  $5x + 2y - 8 = 0$ .

Substituting P in above line,

$$5x + 2y - 8 = 5 \cdot \frac{4}{3} + 2 \cdot \frac{2}{3} - 8 = \frac{20 + 4 - 24}{3} = 0$$

P lies on the third line  $5x + 2y - 8 = 0$

$\therefore$  The given lines are concurrent.

5. Find the distances between the following pairs of parallel straight lines :

i).  $9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$

**Sol.** Given equation is

$$9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0.$$

From above equation  $a = 9, b = 1, c = 8, h = -3, g = 9, f = -3$ .

$$\text{Distance between parallel lines} = 2 \sqrt{\frac{g^2 - ac}{a(a+b)}}$$

$$= 2\sqrt{\frac{9^2 - 9.8}{9(9+1)}} = 2\sqrt{\frac{9}{9.10}} = \sqrt{\frac{4}{10}} = \sqrt{\frac{2}{5}}$$

ii.  $x^2 + 2\sqrt{3}xy + 3y^2 - 3x - 3\sqrt{3}y - 4 = 0$

ans.  $\frac{5}{2}$

6. Show that the pairs of lines  $3x^2 + 8xy - 3y^2 = 0$  and  $3x^2 + 8xy - 3y^2 + 2x - 4y - 1 = 0$  form a squares.

Sol. Equation of the first pair of lines is  $3x^2 + 8xy - 3y^2 = 0$   
 $\Rightarrow (x + 3y)(3x - y) = 0 \Rightarrow 3x - y = 0, x + 3y = 0$

Equations of the lines are  $3x - y = 0$  .....(1) and  $x + 3y = 0$  .....(2)

Equation of the second pair of lines is  $3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = 0$

Since  $3x^2 + 8xy - 3y^2 = (x + 3y)(3x - y)$

Let  $3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = (3x - y + c_1)(x + 3y + c_2)$

Equating the co-efficient of x, we get  $c_1 + 3c_2 = 2$

Equation the co-efficient of y, we get  $3c_1 + c_2 = -4$

$$\begin{array}{ccc} c_1 & c_2 & 1 \\ 3 & 2 & 1 \\ -1 & 4 & -3 \end{array}$$

$$\frac{c_1}{12-2} = \frac{c_2}{-6-4} = \frac{1}{-1-9}$$

$$c_1 = \frac{10}{-10} = -1, c_2 = \frac{-10}{-10} = 1$$

Equations of the lines represented by  $3x^2 + 8xy - 3y^2 + 2x - 4y + 1 = 0$  are

$$3x - y - 1 = 0 \dots(3) \text{ and } x + 3y + 1 = 0 \dots(4)$$

From above equations, lines (1) and (3) are parallel and lines (2) and (4) are parallel.

Therefore given lines form a parallelogram.

But the adjacent sides are perpendicular, it is a rectangle. (since, (1), (2) are perpendicular and (3), (4) are perpendicular.)

The point of intersection of the pair of lines  $3x^2 + 8xy - 3y^2 = 0$  is  $O(0,0)$ .

$$\text{Length of the perpendicular from } O \text{ to (3)} = \frac{|0+0+1|}{\sqrt{1+9}} = \frac{1}{\sqrt{10}}$$

$$\text{Length of the perpendicular from } O \text{ to (4)} = \frac{|0+0+1|}{\sqrt{1+9}} = \frac{1}{\sqrt{10}}$$

Therefore,  $O$  is equidistant from lines (3), (4).

Therefore, the distance between the parallel lines is same. Hence the rectangle is a square.

### III

1. Find the product of the length of the perpendiculars drawn from  $(2,1)$  upon the lines  $12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = 0$

**Sol.** Given pair of lines is  $12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = 0$

Now

$$\begin{aligned} 12x^2 + 25xy + 12y^2 &= 12x^2 + 16xy + 9xy + 12y^2 \\ &= 4x(3x + 4y) + 3y(3x + 4y) = (3x + 4y)(4x + 3y) \end{aligned}$$

$$\text{Let } 12x^2 + 25xy + 12y^2 + 10x + 11y + 2 = (3x + 4y + c_1)(4x + 3y + c_2)$$

Equating the co-efficient of  $x$ ,  $y$  we get

$$4c_1 + 3c_2 = 10 \Rightarrow 4c_1 + 3c_2 - 10 = 0 \dots(1)$$

$$3c_1 + 4c_2 = 11 \Rightarrow 3c_1 + 4c_2 - 11 = 0 \dots(2)$$

Solving,

$$\begin{array}{ccc} c_1 & c_2 & 1 \\ 3 & -10 & 4 \\ 4 & -11 & 3 \end{array}$$

$$\frac{c_1}{-33+40} = \frac{c_2}{-30+44} = \frac{1}{16-9}$$

$$c_1 = \frac{7}{7} = 1, c_2 = \frac{14}{7} = 2$$


Therefore given lines are  $3x + 4y + 1 = 0$  -----(3) and  $4x + 3y + 2 = 0$  -----(4)

$$\text{Length of the perpendicular from } P(2,1) \text{ on (1)} = \frac{6+4+1}{\sqrt{9+16}} = \frac{11}{5}$$

$$\text{Length of the perpendicular from } P(2,1) \text{ on (2)} = \frac{|8+3+2|}{\sqrt{16+9}} = \frac{13}{5}$$

$$\text{Product of the length of the perpendicular} = \frac{11}{5} \times \frac{13}{5} = \frac{143}{25}$$

2. Show that the straight lines  $y^2 - 4y + 3 = 0$  and  $x^2 + 4xy + 4y^2 + 5x + 10y + 4 = 0$  form a parallelogram and find the lengths of its sides.



**Sol.** Equation of the first pair of lines is

$$y^2 - 4y + 3 = 0, \Rightarrow (y-1)(y-3) = 0$$

$$\Rightarrow y - 1 = 0 \text{ or } y - 3 = 0$$

$$\Rightarrow \text{Equations of the lines are } y - 1 = 0 \text{ .....(1)}$$

$$\text{and } y - 3 = 0 \text{ .....(2)}$$

Equations of (1) and (2) are parallel.

$$\text{Equation of the second pair of lines is } x^2 + 4xy + 4y^2 + 5x + 10y + 4 = 0$$

$$\Rightarrow (x + 2y)^2 + 5(x + 2y) + 4 = 0$$

$$\Rightarrow (x + 2y)^2 + 4(x + 2y) + (x + 2y) + 4 = 0$$


$$\Rightarrow (x + 2y)(x + 2y + 4) + 1(x + 2y + 4) = 0$$

$$\Rightarrow (x + 2y + 1)(x + 2y + 4) = 0$$

$$\Rightarrow x + 2y + 1 = 0, x + 2y + 4 = 0$$

$$\text{Equations of the lines are } x + 2y + 1 = 0 \text{ .....(3) and } x + 2y + 4 = 0 \text{ .....(4)}$$

Equations of (3) and (4) are parallel .



$$\text{Solving (1), (3) } x + 2 + 1 = 0, x = -3$$

Co-ordinates of A are (-3, 1)

$$\text{Solving (2), (3) } x + 6 + 1 = 0, x = -7$$

Co-ordinates of D are (-7,3)

Solving (1), (4)  $x + 2 + 4 = 0$ ,  $x = -6$

Co-ordinates of B are (-6, 1)

$$AB = \sqrt{(-3+6)^2 + (1-1)^2} = \sqrt{9+0} = 3$$

$$AD = \sqrt{(-3+7)^2 + (1-3)^2} = \sqrt{16+4} = \sqrt{20} = 2\sqrt{5}$$

Length of the sides of the parallelogram are  $3, 2\sqrt{5}$

3. **Show that the product of the perpendicular distances from the origin to the pair of straight lines represented by  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  is**

$$\frac{|c|}{\sqrt{(a-b)^2 + 4h^2}}$$

**Sol.** Let  $l_1x + m_1y + n_1 = 0$  .....(1)

$l_2x + m_2y + n_2 = 0$  .....(2) be the lines represented by

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$

$$\Rightarrow ax^2 + 2hxy + by^2 + 2gx + 2fy + c = (l_1x + m_1y + n_1)(l_2x + m_2y + n_2)$$

$$\Rightarrow l_1l_2 = a, m_1m_2 = b, l_1m_2 + l_2m_1 = 2h$$

$$l_1n_2 + l_2n_1 = 2g, m_1n_2 + m_2n_1 = 2f, n_1n_2 = c$$

$$\text{Perpendicular from origin to (1)} = \frac{|n_1|}{\sqrt{l_1^2 + m_1^2}}$$

$$\text{Perpendicular from origin to (2)} = \frac{|n_2|}{\sqrt{l_2^2 + m_2^2}}$$

Product of perpendiculars



$$\begin{aligned}
&= \frac{|n_1|}{\sqrt{l_1^2 + m_1^2}} \cdot \frac{|n_2|}{\sqrt{l_2^2 + m_2^2}} \\
&= \frac{|n_1 n_2|}{\sqrt{l_1^2 l_2^2 + m_1^2 m_2^2 + l_1^2 m_2^2 + l_2^2 m_1^2}} \\
&= \frac{|n_1 n_2|}{\sqrt{(l_1 l_2 - m_1 m_2)^2 + (l_1 m_2 + l_2 m_1)^2}} \\
&= \frac{|c|}{\sqrt{(a-b)^2 + (2h)^2}} = \frac{|c|}{\sqrt{(a-b)^2 + 4h^2}}
\end{aligned}$$

4. If the equation  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents a pair of intersecting lines, then show that the square of the distance of their point of intersection from the origin is  $\frac{c(a+b) - f^2 - g^2}{ab - h^2}$ . Also show that the square of this distance from origin is  $\frac{f^2 + g^2}{h^2 + b^2}$  if the given lines are perpendicular.

**Sol.** Let  $l_1x + m_1y + n_1 = 0$  .....(1)

$l_2x + m_2y + n_2 = 0$  .....(2)

be the lines represented by  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

$\Rightarrow ax^2 + 2hxy + by^2 + 2gx + 2fy + c$

$= (l_1x + m_1y + n_1)(l_2x + m_2y + n_2)$

$l_1l_2 = a, m_1m_2 = b, l_1m_2 + l_2m_1 = 2h$

$l_1n_2 + l_2n_1 = 2g, m_1n_2 + m_2n_1 = 2f, n_1n_2 = c$  **Solving (1) and (2)**

$$\frac{x}{m_1n_2 - m_2n_1} = \frac{y}{l_2n_1 - l_1n_2} = \frac{1}{l_1m_2 - l_2m_1}$$

The point of intersection is  $P = \left[ \frac{m_1 n_2 - m_2 n_1}{l_1 m_2 - l_2 m_1}, \frac{l_2 n_1 - l_1 n_2}{l_1 m_2 - l_2 m_1} \right]$

$$\begin{aligned} OP^2 &= \frac{(m_1 n_2 - m_2 n_1)^2 + (l_2 n_1 - l_1 n_2)^2}{(l_1 m_2 - l_2 m_1)^2} \\ &= \frac{(m_1 n_2 + m_2 n_1)^2 - 4m_1 m_2 n_1 n_2 + (l_1 n_2 + l_2 n_1)^2 - 4l_1 l_2 n_1 n_2}{(l_1 m_2 + l_2 m_1)^2 - 4l_1 l_2 m_1 m_2} \\ &= \frac{4f^2 - 4abc + 4g^2 - 4ac}{4h^2 - 4ab} \\ &= \frac{c(a+b) - f^2 - g^2}{ab - h^2}. \end{aligned}$$

If the given pair of lines are perpendicular, then  $a + b = 0 \Rightarrow a = -b$


$$\Rightarrow OP^2 = \frac{0 - f^2 - g^2}{(-b)b - h^2} = \frac{f^2 + g^2}{h^2 + b^2}$$

## HOMOGENISATION

### THEOREM

The equation to the pair of lines joining the origin to the points of intersection of the curve  $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  and the line

$$L \equiv lx + my + n = 0 \text{ is } ax^2 + 2hxy + by^2 + (2gx + 2fy) \left( \frac{lx + my}{-n} \right) + c \left( \frac{lx + my}{-n} \right)^2 = 0 \text{ ---(1)}$$



Eq (1) represents the combined equation of the pair of lines  $\overline{OA}$  and  $\overline{OB}$ .

**EXERCISE - 4(c)**

**I**

- 1. Find the equation of the lines joining the origin to the point of intersection of  $x^2 + y^2 = 1$  and  $x + y = 1$**

**Sol.** The given curves are  $x^2 + y^2 = 1$  .....(1)

$$x + y = 1 \quad \text{.....(2)}$$

Homogenising (1) with the help of (2) then  $x^2 + y^2 = 1^2$

$$\Rightarrow x^2 + y^2 = (x + y)^2 = x^2 + y^2 + 2xy \text{ i.e. } 2xy = 0 \Rightarrow xy = 0$$

- 2. Find the angle between the lines joining the origin to points of intersection of  $y^2 = x$  and  $x + y = 1$ .**

**Sol.** Equation of the curve is  $y^2 = x$  .....(1) and Equation of line is  $x + y = 1$  .....(2)

Harmogonsing (1) with the help of (2)

$$Y^2 -x.1 =0 \Rightarrow y^2 = x(x + y) = x^2 + xy$$


$\Rightarrow x^2 + xy - y^2 = 0$  which represents a pair of lines. From this equation

$$a + b = 1 - 1 = 0$$

The angle between the lines is  $90^0$ .

**II**

- 1. Show that the lines joining the origin to the points of intersection of the curve  $x^2 - xy + y^2 + 3x + 3y - 2 = 0$  and the straight line  $x - y - \sqrt{2} = 0$  are mutually perpendicular.**



Sol.

Let A, B be the points of intersection of the line and the curve.

$$\text{Equation of the curve is } x^2 - xy + y^2 + 3x + 3y - 2 = 0 \dots\dots(1)$$

$$\text{Equation of the line AB is } x - y - \sqrt{2} = 0$$

$$\Rightarrow x - y = \sqrt{2} \Rightarrow \frac{x - y}{\sqrt{2}} = 1 \dots\dots(2)$$

Homogenising, (1) with the help of (2) combined equation of OA, OB is

$$x^2 - xy + y^2 + 3x \cdot 1 + 3y \cdot 1 - 2 \cdot 1^2 = 0$$

$$\Rightarrow x^2 - xy + y^2 + 3(x + y) \frac{x - y}{\sqrt{2}} - 2 \frac{(x - y)^2}{2} = 0$$

$$\Rightarrow x^2 - xy + y^2 + \frac{3}{\sqrt{2}}(x^2 - y^2) - (x^2 - 2xy + y^2) = 0$$

$$\Rightarrow x^2 - xy + y^2 + \frac{3}{\sqrt{2}}x^2 - \frac{3}{\sqrt{2}}y^2 - x^2 + 2xy - y^2 = 0$$

$$\Rightarrow \frac{3}{\sqrt{2}}x^2 + xy - \frac{3}{\sqrt{2}}y^2 = 0$$


$$\Rightarrow a + b = \frac{3}{\sqrt{2}} - \frac{3}{\sqrt{2}} = 0$$

$\therefore$  OA, OB are perpendicular.

2. Find the values of  $k$ , if the lines joining the origin to the points of intersection of the curve  $2x^2 - 2xy + 3y^2 + 2x - y - 1 = 0$  and the line  $x + 2y = k$  are mutually perpendicular.

**Sol.** Given equation of the curve is  $S \equiv 2x^2 - 2xy + 3y^2 + 2x - y - 1 = 0 \dots\dots(1)$

Equation of AB is  $x + 2y = k$



$$\frac{x + 2y}{k} = 1 \dots\dots(2)$$

Let A, B be the points of intersection of the line and the curve.

Homogenising, (1) with the help of (2), the combined equation of OA, OB is

$$2x^2 - 2xy + 3y^2 + 2x \cdot 1 - y \cdot 1 - 1^2 = 0$$

$$2x^2 - 2xy + 3y^2 + 2x \frac{(x + 2y)}{k} - y \frac{(x + 2y)}{k} - \frac{(x + 2y)^2}{k^2} = 0$$

$$\Rightarrow 2k^2x^2 - 2k^2xy + 3k^2y^2 + 2kx(x + 2y) - ky(x + 2y) - (x + 2y)^2 = 0$$

$$\Rightarrow 2k^2x^2 - 2k^2xy + 3k^2y^2 + 2kx^2 + 4kxy - kxy - 2ky^2 - x^2 - 4xy - 4y^2 = 0$$

$$\Rightarrow (2k^2 + 2k - 1)x^2 + (-2k^2 + 3k - 4)xy + (3k^2 - 2k - 4)y^2 = 0$$


Given that above lines are perpendicular, Co-efficient  $x^2$  + co-efficient of  $y^2 = 0$

$$\Rightarrow 2k^2 + 2k - 1 + 3k^2 - 2k - 4 = 0$$

$$\Rightarrow 5k^2 = 5 \Rightarrow k^2 = 1 \therefore k = \pm 1$$

3. Find the angle between the lines joining the origin to the points of intersection of the curve  $x^2 + 2xy + y^2 + 2x + 2y - 5 = 0$  and the line  $3x - y + 1 = 0$

Sol.



Equation of the curve is  $x^2 + 2xy + y^2 + 2x + 2y - 5 = 0 \dots\dots(1)$

Equation of AB is  $3x - y + 1 = 0 \Rightarrow y - 3x = 1 \dots\dots(2)$

Let A,B the the points of intersection of the line and the curve.

Homogenising (1) with the help of (2), combined equation of OA, OB is

$$x^2 + 2xy + y^2 + 2x \cdot 1 + 2y \cdot 1 - 5 \cdot 1^2 = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + 2x(y - 3x) + 2y(y - 3x) - 5(y - 3x)^2 = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + 2xy - 6x^2 + 2y^2 - 6xy - 5(y^2 + 9x^2 - 6xy) = 0$$

$$\Rightarrow -5x^2 - 2xy + 3y^2 - 5y^2 - 45x^2 + 30xy = 0$$

$$\Rightarrow -50x^2 + 28xy - 2y^2 = 0 \Rightarrow 25x^2 - 14xy + y^2 = 0$$


let  $\theta$  be the angle between OA and OB ,then

$$\begin{aligned} \cos \theta &= \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}} = \frac{|25+1|}{\sqrt{(25-1)^2 + 196}} = \frac{26}{\sqrt{576+196}} = \frac{26}{\sqrt{772}} \\ &= \frac{26}{2\sqrt{193}} = \frac{13}{\sqrt{193}} \therefore \theta = \cos^{-1}\left(\frac{13}{\sqrt{193}}\right) \end{aligned}$$

### III

1. Find the condition for the chord  $lx + my = 1$  of the circle  $x^2 + y^2 = a^2$  (whose centre is the origin) to subtend a right angle at the origin.

Sol.



Equation of the circle  $x^2 + y^2 = a^2$  .....(1)

Equation of AB is  $lx + my = 1$  .....(2)

Let A,B the the points of intersection of the line and the curve

Homogenising (1) with the help of (2), the combined equation of OA, OB is

$$x^2 + y^2 = a^2 \cdot 1^2 \Rightarrow x^2 + y^2 = a^2 (lx + my)^2$$

$$= a^2 (l^2 x^2 + m^2 y^2 + 2lmxy) = a^2 l^2 x^2 + a^2 m^2 y^2 + 2a^2 lmxy$$

$$\Rightarrow a^2 l^2 x^2 + 2a^2 lmxy + a^2 m^2 y^2 - x^2 - y^2 = 0$$


$$\Rightarrow (a^2 l^2 - 1)x^2 + 2a^2 lmxy + (a^2 m^2 - 1)y^2 = 0$$

Since OA, OB are perpendicular, Coefficient of  $x^2$  + co-efficient of  $y^2 = 0$

$$\Rightarrow a^2 l^2 - 1 + a^2 m^2 - 1 = 0 \Rightarrow a^2 (l^2 + m^2) = 2 \text{ which is the required condition}$$

2. Find the condition for the lines joining the origin to the points of intersection of the circle  $x^2 + y^2 = a^2$  and the line  $lx + my = 1$  to coincide.

Sol.



Equation of the circle is  $x^2 + y^2 = a^2$  .....(1)

Equation of AB is  $lx + my = 1$  .....(2) .

Let A, B the the points of intersection of the line and the curve.

Homogenising (1) with the help of (2) ,

Then the combined equation of OA, OB is  $x^2 + y^2 = a^2 \cdot l^2$

$$x^2 + y^2 = a^2 (lx + my)^2 = a^2 (l^2x^2 + m^2y^2 + 2lmxy)$$

$$\Rightarrow x^2 + y^2 = a^2 l^2 x^2 + a^2 m^2 y^2 + 2a^2 lmxy$$

$$\Rightarrow (a^2 l^2 - 1)x^2 + 2a^2 lmxy + (a^2 m^2 - 1)y^2 = 0$$

Since OA, OB are coincide  $\Rightarrow h^2 = ab$

$$\Rightarrow a^4 l^2 m^2 = (a^2 l^2 - 1)(a^2 m^2 - 1) \Rightarrow a^4 l^2 m^2 = a^4 l^2 m^2 - a^2 l^2 - a^2 m^2 + 1$$

$$\therefore a^2 l^2 - a^2 m^2 + 1 = 0 \Rightarrow a^2 (l^2 + m^2) = 1$$

This is the required condition.



3. Write down the equation of the pair of straight lines joining the origin to the points of intersection of the lines  $6x - y + 8 = 0$  with the pair of straight lines  $3x^2 + 4xy - 4y^2 - 11x + 2y + 6 = 0$ . Show that the lines so obtained make equal angles with the coordinate axes.

**Sol.** Given pair of line is  $3x^2 + 4xy - 4y^2 - 11x + 2y + 6 = 0 \dots(1)$

$$\text{Given line is } 6x - y + 8 = 0 \Rightarrow \frac{6x - y}{-8} = 1 \Rightarrow \frac{y - 6x}{8} = 1 \dots\dots(2)$$

Homogenising (1) w.r.t (2)

$$3x^2 + 4xy - 4y^2 - (11x - 2y)\left(\frac{y - 6x}{8}\right) + 6\left(\frac{y - 6x}{8}\right)^2 = 0$$

$$64[3x^2 + 4xy - 4y^2] - 8[11xy - 66x^2 - 2y^2 + 12xy] + 6[y^2 + 36x^2 - 12xy] = 0$$

$$\Rightarrow 936x^2 + 256xy - 256xy - 234y^2 = 0$$

$$\Rightarrow 468x^2 - 117y^2 = 0$$

$$\Rightarrow 4x^2 - y^2 = 0 \dots\dots (3)$$

Is eq. of pair of lines joining the origin to the point of intersection of (1) and (2).

$$\text{The eq. pair of angle bisectors of (3) is } h(x^2 - y^2) - (a - b)xy = 0$$

$$\Rightarrow 0(x^2 - y^2) - (4 - 1)xy = 0 \Rightarrow xy = 0$$

$x = 0$  or  $y = 0$  which are the eqs. is of co-ordinates axes

$\therefore$  The pair of lines are equally inclined to the co-ordinate axes

4. **If the straight lines given by  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  intersect on Y-axis, show that  $2fgh - bg^2 - ch^2 = 0$**

**Sol.** Given pair of lines is  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

Equation of Y-axis is  $x = 0$  then equation becomes  $by^2 + 2fy + c = 0 \dots\dots(1)$

Since the given pair of lines intersect on Y – axis, the roots of equation (1) are equal.

$\therefore$  Discriminate = 0

$$\Rightarrow (2f)^2 - 4.b.c = 0 \Rightarrow 4f^2 - 4bc = 0$$

$$\Rightarrow f^2 - bc = 0 \Rightarrow f^2 = bc$$

Since the given equation represents a pair of lines

$$abc + 2fgh + af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow a(f^2) + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow 2fgh - bg^2 - ch^2 = 0$$

5. **Prove that the lines represented by the equations  $x^2 - 4xy + y^2 = 0$  and  $x + y = 3$  form an equilateral triangle.**


**Sol.** Since the straight line  $L : x + y = 3$  makes  $45^\circ$  with the negative direction of the

X –axis, none of the lines which makes  $60^\circ$  with the line L is vertical. If ‘m’ is the

slope of one such straight line, then  $\sqrt{3} = \tan 60^\circ = \left| \frac{m+1}{1-m} \right|$  and so, satisfies the

$$\text{equation } (m+1)^2 = 3(m-1)^2$$

$$\text{Or } m^2 - 4m + 1 = 0 \dots\dots\dots(1)$$



But the straight line having slope 'm' and passing through the origin is

$$y = mx \dots\dots\dots (2)$$

So the equation of the pair of lines passing through the origin and inclined at  $60^\circ$  with the line L is obtained by eliminating 'm' from the equations (1) and (2).

Therefore the combined equation of this pair of lines is  $\left(\frac{y}{x}\right)^2 - 4\left(\frac{y}{x}\right) + 1 = 0$  (i.e.)

$$x^2 - 4xy + y^2 = 0$$

Which is the same as the given pair of lines. Hence, the given traid of lines form an equilateral triangle.

6. Show that the product of the perpendicular distances from a point  $(\alpha, \beta)$  to

the pair of straight lines  $ax^2 + 2hxy + by^2 = 0$  is  $\frac{|a\alpha^2 + 2h\alpha\beta + b\beta^2|}{\sqrt{(a-b)^2 + 4h^2}}$

Sol. Let  $ax^2 + 2hxy + by^2 \equiv (l_1x + m_1y)(l_2x + m_2y)$

Then the separate equations of the lines represented by the equation

$$ax^2 + 2hxy + by^2 = 0 \text{ are } L_1 : l_1x + m_1y = 0 \text{ and } L_2 : l_2x + m_2y = 0$$

Also, we have  $l_1l_2 = a; m_1m_2 = b$  and  $l_1m_2 + l_2m_1 = 2h$

$$d_1 = \text{length of the perpendicular from } (\alpha, \beta) \text{ to } L_1 = \frac{|l_1\alpha + m_1\beta|}{\sqrt{l_1^2 + m_1^2}}$$

$$d_2 = \text{length of the perpendicular from } (\alpha, \beta) \text{ to } L_2 = \frac{|l_2\alpha + m_2\beta|}{\sqrt{l_2^2 + m_2^2}}$$

Then, the product of the lengths of the perpendiculars from  $(\alpha, \beta)$  to the given pair of lines  $= d_1 d_2$

$$= \frac{|(l_1 \alpha + m_1 \beta)(l_2 \alpha + m_2 \beta)|}{\sqrt{(l_1^2 + m_1^2)(l_2^2 + m_2^2)}} = \frac{|a\alpha^2 + 2h\alpha\beta + b\beta^2|}{\sqrt{(a-b)^2 + 4h^2}}$$

### PROBLEMS FOR PRACTICE.

1. If the lines  $xy + x + y + 1 = 0$  and  $x + ay - 3 = 0$  are concurrent, find  $a$ .
2. The equation  $ax^2 + 3xy - 2y^2 - 5x + 5y + c = 0$  represents two straight lines perpendicular to each other. Find  $a$  and  $c$ .
3. Find  $\lambda$  so that  $x^2 + 5xy + 4y^2 + 3x + 2y + \lambda = 0$  may represent a pair of straight lines. Find also the angle between them for this value of  $\lambda$ .
4. If  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  represents the straight lines equidistant from the origin, show that  $f^4 - g^4 = c(bf^2 - ag^2)$
5. Find the centroid of the triangle formed by the lines  $12x^2 - 20xy + 7y^2 = 0$  and  $2x - 3y + 4 = 0$

ANS.  $= \left( \frac{8}{3}, \frac{8}{3} \right)$

6. Let  $aX^2 + 2hXY + bY^2 = 0$  represent a pair of straight lines. Then show that the equation of the pair of straight lines.

i) Passing through  $(x_o, y_o)$  and parallel to the given pair of lines is

$a(x - x_o)^2 + 2h(x - x_o)(y - y_o) + b(y - y_o)^2 = 0$  ii) Passing through  $(x_o, y_o)$  and perpendicular to the given pair of lines

is  $b(x - x_o)^2 - 2h(x - x_o)(y - y_o) + a(y - y_o)^2 = 0$

7. Find the angle between the straight lines represented by  
 $2x^2 + 3xy - 2y^2 - 5x + 5y - 3 = 0$
8. Find the equation of the pair of lines passing through the origin and perpendicular to the pair of lines  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$
9. If  $x^2 + xy + 2y^2 + 4x - y + k = 0$  represents a pair of straight lines find k.
10. Prove that equation  $2x^2 + xy - 6y^2 + 7y - 2 = 0$  represents a pair of straight line.
11. Prove that the equation  $2x^2 + 3xy - 2y^2 - x + 3y - 1 = 0$  represents a pair of perpendicular straight lines.
12. Show that the equation  $2x^2 - 13xy - 7y^2 + x + 23y - 6 = 0$  represents a pair of straight lines. Also find the angle between the co-ordinates of the point of intersection of the lines.
13. Find that value of  $\lambda$  for which the equation  
 $\lambda x^2 - 10xy + 12y^2 + 5x - 16y - 3 = 0$  represents a pair of straight lines.
14. Show that the pair of straight lines  $6x^2 - 5xy - 6y^2 = 0$  and  $6x^2 - 5xy - 6y^2 + x + 5y - 1 = 0$  form a square.
15. Show that the equation  $8x^2 - 24xy + 18y^2 - 6x + 9y - 5 = 0$  represents pair of parallel straight lines are find the distance between them.